

A study of pollen viability and longevity in Gokarn (*Clitoria ternatea*) a medicinal plant

S.R. KARAD, A.S. JAGTAP, A.U. KUBADE AND P.D.KHADE

Accepted : March, 2009

SUMMARY

Clitoria ternatea L, a naturally grown medicinal plant, endophyte free, planted in Dhanvantari Udyan, College of Agriculture, Kolhapur was selected for present investigation. Locally it is called as Gokarn / Asian pigeonwing, used to study the viability of pollen grains and its longevity. Among different staining methods, tetrazolium test used to assess pollen viability. The fully fertile, five plants randomly selected for collection of pollens from 7 to 11.30 a.m. at every half an hour interval. These pollen grains were stained by using 0.3% TTC (2,3,5-triphenyl tetrazolium chloride) prepared in 0.5% and 1% sucrose solution to prevent bursting and swelling of pollen grains and incubated in dark at $30\pm 2^\circ\text{C}$ for 30 and 60 min, respectively at 70 % relative humidity. At the end of incubation period, viable pollen grains turned red and central area was observed. The more viability is observed in 30 min incubation period in both 0.5 and 1% sucrose solution as compared with 60 min incubation period. In 0.5% sucrose solution, the more viability was observed between 8-10 a.m. and the viability was less before 8 and after 10 a.m. in 30 min incubation. Similar results were observed in 1 % sucrose solution in 30 min incubation. In 60 min incubation period, the pollens were more viable between 8-10.30 a.m. and less viable before 8 and after 10.30 a.m. in 0.5% sucrose solution. While in same incubation period (60 min) with 1 % sucrose solution showed more viability between 8-10 a.m. and less viability before 8 and after 10 a.m. From these observations, it has been concluded that the pollen viability and longevity of *Clitoria ternatea* L. was maximum at 8-10 a.m. during October 2007 at Kolhapur centre. These properties of pollen should be considered to increase efficiency of breeding programme and selection of suitable pollinizer in *Clitoria ternatea* L. a important medicinal plant.

Key words : Gokarn, Pollen, Medicinal plant

The study of pollen viability after dehiscence is crucial for successful pollination (Stone *et al.*, 1995). The pollen viability determines the possibility of outcrossing between species. It also provides limited specific reproductive barriers. The important medicinal plant Gokarn (*Clitoria ternatea*) was used throughout the study. It is imperative to know the extent of viability of pollen sample to use for experiment and pollination. Its viability needs to monitor under different storage condition. The objective of study was to optimise the availability and viability of gokarn pollen. The viability, tube growth and morphological homogeneity is related to pollen quality is important factor in Gokarn. These properties are important for plant breeder, geneticist and growers for selection of suitable pollen and hybridization method. This study of pollen viability and longevity of Gokarn is important in evaluating outcrossing frequencies. Therefore, to determine the actual amount of viable pollen and to find out viability of pollen grains to optimize viability and inter

relationship between them the present study was undertaken.

MATERIALS AND METHODS

The important naturally grown cultivar of gokarn (*Clitoria ternatea*) plants was used in the present investigation. The plants for experiment were planted in Dhanvantari garden, College of Agriculture, Kolhapur. A number of staining methods were used for testing pollen viability. Among these methods tetrazolium test was used in this study. In many species tetrazolium test (TZ) has proved satisfactory in assessing pollen viability. The pollen grains showed a gradation in colour development, hence judging a cut off points for colour intensity to determine viability became subjective. The five fully fertile plants are randomly selected for collection of pollens. The pollen grains were collected from 7 a. m. to 11.30 a.m. at an interval of every half an hour. The collected pollens were stained by using 0.3% TTC (2, 3, 5-triphenyl tetrazolium chloride) solution. The TTC solution was prepared in 0.5% and 1 % sucrose solution to prevent bursting of pollen grains. A small amount of pollens suspended on microslide with TTC drop and coverslip was applied. Then the microslide was transferred to humidity chamber and incubated in the dark $30\pm 2^\circ\text{C}$ for 30 min and 60 min,

Correspondence to:

S.R. KARAD, Department of Agricultural Botany, College of Agriculture, KOLHAPUR (M.S.) INDIA

Authors' affiliations:

A.S. JAGTAP, A.U. KUBADE AND P.D. KHADE, Department of Agricultural Botany, College of Agriculture, KOLHAPUR (M.S.)

respectively. At the end of incubation period, the slides were observed under trinocular research microscope and scored for percentage of viable pollen grains *i.e.* pollen grains that have turned red due to accumulation of formazan, only central area was noted as suggested by Shivanna and Rangaswamy.

RESULTS AND DISCUSSION

The successful germination of pollen is most immediately after anthesis and viability deteriorates rapidly in most species (Kearns and Inouye, 1993). The pollen viability decreases significantly with an hour of exposure to air after dehiscence. However, not all pollens are short lived; some roseaceae and liliaceae pollen can remain viable for 100 days. (Leduc *et al.*, 1990; Song, 2001). In present investigation, results confirmed the rapid pollen viability in Gokarn. In natural condition, pollen grains were viable in a day between 7 to 11.30 a.m. In present study, pollen grains were stored under $652 \pm 2\%$ humidity for 30 min and 60 min incubation period. But more viability was observed in 30 min incubation period in both 0.5 % and 1 % sucrose solution, compared to 60 min incubation period (Table 1) (Mark and David, 2003).

In 0.5% sucrose solution, the more viability was observed between 8-10 a.m. with 30 min incubation. While the viability was less before 8 a.m. and after 10 a.m. Similar results were observed in 1% sucrose solution.

In 60 min incubation period, the pollens were more viable between 8-10.30 a.m. and less viable before 8 a.m. and after 10.30 a.m. in 0.5% sucrose solution. While same incubation period with 1% sucrose solution showed more viability between 8-10 am and less viability before 8 am

Table 1 : Pollen viability score in different sucrose solution of Gokarn plant

Pollen collection time at (a.m.)	0.5 % sucrose solution		1 % sucrose solution	
	30 min	60 min	30 min	60 min
7.30	2	1	3	2
8.00	4	3	4	3
8.30	4	3	4	3
9.00	3	3	4	3
9.30	4	3	3	3
10.00	4	3	4	3
10.30	3	3	3	2
11.00	2	2	2	3
11.30	2	2	3	3

Score grade: 1. No viable, 2. Less viable, 3. Moderate viable, 4. Yiable, 5. Fully viable

and after 10 a.m. (Rodriguez *et al.*, 2000)

From the results, it has been concluded that the pollen viability and longevity of gokarn (*Clitoria ternatea*) was maximum between 8-10 a.m. at Kolhapur location in the month October 2007. Hence, these properties of pollen should be considered to increase efficiency of breeding programme and selection of suitable pollinizer in medicinal plant.

Acknowledgement:

The authors are thankful to Dr. D.M. Sawant, Associate Dean, College of Agriculture, Kolhapur for providing the facilities and grateful to C.E.O., NMPB, New Delhi for financial assistance for exploration and inventorization of medicinal plants and its research trials under adhoc projects.

REFERENCES

- Kearns, C.A. and Inouye, D.W. (1993). *Techniques for pollination biologists*. University Press of Colorado, 583.
- Leduc, N., Monnier, M. and Douglas, G.C. (1990). Germination of trinucleated pollen: Formulation of new medium for *Capsilla bursa-pastoris*. *Sexual Plant Reproduction*, **3** : 228-235.
- Mark, Gibernau and David, Macquart (2003). Pollen viability and longevity in two species of *Arum*. *Aroideana*, **26** : 58-62.
- Rodriguez-Riano and Dafui, T.A. (2000). A new procedure to pollen viability. *Sexual Plant Reproduction*, **12** : 241-244.
- Shivanna, K.R. and Rangaswamy, M.S. Lab manual of pollen biology. pp-33-37.
- Song, Z.P. (2001). A study of pollen viability and longevity in *Oryza rufipogon*, *O. sativa*, and their hybrids. *Genetic Res.*, **26** (2) : 31-34.
- Stone, J.L., Thomson, J.D. and Dent-Acosta, S.J. (1995). Assesement of pollen viability in hand pollination experiments: a review. *American J. Bot.*, **82** : 1186-1197.

