

Antioxidant activity and phenolic contents in *Ocimum sanctum* and *Ocimum basilicum*

INAMPUDI SAILAJA¹, IVVALA ANAND SHAKER² AND YEDLAPALLI KEERTHI RATNA¹

¹Department of Biochemistry and Molecular Biology, College of Agricultural Biotechnology, Loni, AHMEDNAGAR, (M.S.) INDIA

²Department of Biochemistry, Rural Medical College, P.I.M.S, Loni, AHMEDNAGAR, (M.S.) INDIA

(Accepted : November, 2009)

Ocimum sanctum Linn, *Ocimum basilicum* Linn, which have been normally used in traditional puja at every Indian Hindu home, was estimated by three different methods: ferric reducing antioxidant power (FRAP) assay, improved ABTS radical cation decolorization assay, and DPPH free radical scavenging assay. Additionally, their total phenolic contents were analyzed by Folin-Ciocalteu micro method. The result showed that *Ocimum basilicum* seemed to be better sources of antioxidant compounds, followed by *Ocimum sanctum*, respectively. Total phenolic content analysis, were significantly related to those of FRAP, ABTS, DPPH.

Key words : Antioxidant status, Total phenolic content, *O.basilicum*, *O.sanctum*

INTRODUCTION

Plants are one of the most important sources of medicines. *Ocimum basilicum* L. (labiateae) is a grassy and annual plant. The leaves of this plant are oval with a sharp tip and the flowers are yellow, white and pink. It is a native plant of Iran, Afghanistan and India (Mirheidar, 1990; Volak and Plantes 1997; Zargary, 1990). This plant is a popular culinary herb and its essential oils have been used for many years in food perfumery and dental products (Suppakul *et al.*, 2003) in Iranian traditional medicine (Dasgupta *et al.*, 2004; Javanmardi *et al.*, 2002). According to the literature, about 45 compounds are found in volatile oil of this plant and the major compounds are linalol, eugenol, methylchavicol, methylcinnamat, linolen, ocimene, pinene, cineol, anethol, estragol, thymol, citral and comphor (Mirheidar, 1990; Wang *et al.*, 2003 and Grayer *et al.*, 1996). Different parts of *Ocimum basilicum* have diuretic (Mirheidar, 1990; Zargary, 1990), antipyretic (Mirheidar, 1990) and antitussive effects (Mirheidar, 1990; Volak and Plantes 1997) and have been used to treat gastritis, stomach-ache, flatulence and constipation (Mirheidar, 1990; Volak and Plantes, 1997; Zargary, 1990; Avesina, 1990; Singh, 1999; Akthar and Muneer, 1989), vomiting and hiccup (Mirheidar, 1990). The seeds and the boiled extract are used for treatment of diarrhea. This plant also has therapeutic effects for nasal polyps (Mirheidar, 1990), upper respiratory tract diseases and dyspnea (Mirheidar, 1990; Volak and Plantes, 1997); inflammation of the urinary tract (Volak and Plantes, 1997) and it has been

used as a bathing solution for treatment of ulcers (Volak and Plantes, 1997; Avesina, 1990) and increase in lactation (Zargary, 1990). Inhalation of the fragrance of this plant is useful to alleviate headache (Mirheidar, 1990) and also it produces a favorable impression on the mental activity (Satoh and Sugawara, 2003). In the previous studies, the anti-inflammatory (Singh, 1999; Singh, 1998) anti-HIV (Yamasaki *et al.*, 1998), antioxidant (Dasgupta, *et al.*, 2004; Jayasinghe *et al.*, 2003; Yun, *et al.*, 2003; Lee and Shibamoti, 2002), antibacterial and antifungal (Suppakul *et al.*, 2003; Edris and Farrag 2003; Opalchenova and Obreshkova, 2003; Lachowicz *et al.*, 1998). In Ayurveda *Tulsi* (*Ocimum sanctum* L.) has been well documented for its therapeutic potentials and described as Dashemani Shwasaharni (antiasthmatic) and antikaphic drugs (Kaphaghna) (Sirkar, 1989). Although the traditional medical practitioners in India have been widely using this medicinal plant for management of various disease conditions from ancient time, not much is known about the mode of action of *Tulsi* and a rational approach to this traditional medical practice with modern system of medicine is also not available. In last few decades several studies have been carried out by Indian scientists and researchers to suggest the role of essential oils and eugenol in therapeutic potentials of *Ocimum sanctum* L. (Sen, 1993; AOAC, 1997). Eugenol is a phenolic compound and major constituent of essential oils extracted from different parts of *Tulsi* plant. The therapeutic potential of *Tulsi* has been established on the basis of several pharmacological studies carried out with eugenol and steam distilled, petroleum ether and benzene

extracts of different parts of *Tulsi* plant (Sen, 1993; Khanna and Bhatia, 2003). Holy basil (*Ocimum sanctum*, *Ocimum basilicum*), available from fresh markets of Shridi, Taluka –Rahata, Dist –Ahmednagar, Maharashtra by three different methods, viz., ferric reducing antioxidant power (FRAP) assay, improved ABTS radical cation decolorization assay and DPPH free radical scavenging assay. The total phenolic contents of the above culinary plants were also estimated and related to their antioxidant capacity results.

MATERIALS AND METHODS

Sample extraction :

Fresh samples of plants were collected in and around Loni. Their moisture contents were analyzed by the method of AOAC (Official method of analysis of AOAC international, 1997). The sample extraction method of (Leong and Shui, 2002) was modified for sample preparation. Edible portion of the fresh sample was homogenized using a blender. A mixture of deionized water and 95% ethanol was used as extraction solvent and 95% ethanol was selected for preliminary studies. From preliminary studies, 60% (v/v) of 95% ethanol was selected as extraction solvent for chillies and holy basil, while deionized water and 95% ethanol were chosen for garlic and pumpkin, respectively. Two grams of homogenized sample were added with 10 ml of the selected solvent. The extraction was done by using a vortex mixer for 60 seconds. The mixture was filtered and the filtrate was collected and used for FRAP, ABTS, DPPH, and total phenolic content assayed by spectrophotometer.

Ferric reducing antioxidant power (FRAP) assay :

The FRAP was assessed according to Benzie and Strain (1999). Briefly, 6 ml of working FRAP reagent prepared daily was mixed with 20 - 100 μ l of the extract. The absorbance at 593 nm was recorded after 30-min incubation at 37°C. Absorbance increases were calculated as FRAP values by comparing with standard curves created by vitamin C (0 - 15 μ g), and reported as mg vitamin C equivalent per gram of fresh weight.

ABTS radical cation decolorization assay :

The ABTS method of (Re *et al.*, 1999) was modified. ABTS radical cation (ABTS+) was produced by reacting 7 mM ABTS stock solution with 2.45 mM potassium persulphate (final concentration) and allowing the mixture to stand in the dark at room temperature for 12 - 16 hours before use. The ABTS+ solution was diluted with

deionized water and 95 % ethanol (1:1) to an absorbance of 0.70 (+ 0.02) at 734 nm. The extract (20-100 μ l) was mixed with 6 ml of diluted ABTS+ solution. The decrease of absorbance was recorded at 1 min after mixing. Absorbance decreases were calculated as ABTS values by comparing with standard curves created by vitamin C (0 - 20 μ g), and the results were reported as mg vitamin C equivalent per gram of fresh weight.

DPPH free radical scavenging activity :

The method of (Brand-Williams and Cuelier, 1995) was used with some modifications. DPPH radical solution (0.8 mM) in 95% ethanol was prepared. The extract (100-1000 μ l) was diluted to 5.4 ml by deionized water and 95 % ethanol (1:1) before 0.6 ml of the DPPH solution was added and mixed. The decrease of absorbance was recorded at 1 min after mixing. Absorbance decreases were calculated as DPPH values by comparing with standard curves created by vitamin C (0 - 40 μ g), and the results were reported as mg vitamin C equivalent per gram of fresh weight.

Total phenolic content :

The Folin-Ciocalteu micro method of (Waterhouse, 2005) was used to estimate total phenolic content (TPC). The extract (60-300 μ l) was diluted with deionized water to 4.8 ml, and 300 μ l of Folin-Ciocalteu reagent was added and shaken. After 8 min, 900 μ l of 20% sodium carbonate solution was added with mixing. The solution was left at 40°C for 30 min before reading the absorbance at 765 nm. Gallic acid (0 - 50 μ g) was used as standard, and the results were reported as mg gallic acid equivalent per gram of fresh weight.

Calculation and statistical analysis :

The values of FRAP, ABTS, DPPH, and TPC (mg standard equivalent per gram of fresh weight) were calculated using the equation below:

Values of FRAP, ABTS, DPPH, and TPC :

$$(\text{Mg standard equivalent per gram of fresh weight}) = \frac{[(\text{SA} - \text{BA}) / (\text{Slope})][10 / \text{U}]}{[2][1,000]}$$

where,

SA = Sample absorbance for FRAP value and TPC or absorbance decrease of sample for ABTS and DPPH values.

BA = Blank (no extract) absorbance for FRAP value and TPC or absorbance decrease of blank for ABTS and

DPPH values (extract was substituted by demonized water for blank).

- Slope = Slope of standard curve.
- $[10 / U]$ = Total volume of extract (10 ml) / used volume of extract (ml).
- $[2]$ = Weight of used sample (g).
- $[1,000]$ = Factor for changing μg to mg.

Each experiment was performed in triplicate on different purchased samples. Mean comparisons were performed by Duncan's new multiple range test (DMRT). The correlations between all antioxidant status assays and total phenolic contents were analyzed.

RESULTS AND DISCUSSION

The results of the antioxidant capacity assessment of the studied plants as determined by FRAP, ABTS, and DPPH assays are shown in Table 2. These differences could be explained by different mechanisms of analytical methods. FRAP assay measures the ability to reduce a ferric tripyridyltriazine (Fe^{3+} -TPTZ) to a ferrous form (Fe^{2+} -TPTZ) of samples (Leong and Shui, 2002) ABTS and DPPH assays are based on the reduction of ABTS (Re *et al.*, 1999) and DPPH free radicals (Brand-Williams and Cuelier 1995) of samples, but values from DPPH assay might be lower than those from ABTS assay. (Wang and Shao, 1998) showed that some compounds which have ABTS scavenging activity may not show DPPH scavenging activity, and (Arts and Haenen, 2004) found that some products of ABTS scavenging reaction may have a higher antioxidant capacity and can further react with remained ABTS radicals.

Table 2 shows that *O. basilicum* had the highest

Table 1 : Moisture content of plants

Plants	Moisture content (%)
<i>Ocimum sanctum</i>	86.44 + 0.46
<i>Ocimum basilicum</i>	85.90 + 0.61

Table 2 : Antioxidant status and total phenolic content

Plants	FRAP*	ABTS*	DPPH*	TPC**
<i>Ocimum sanctum</i>	3.05 + 0.73 ^a	5.21 + 0.79 ^a	0.42 + 0.22 ^b	1.25 + 0.47 ^b
<i>Ocimum basilicum</i>	1.49 + 0.55 ^b	4.03 + 0.39 ^b	0.52 + 0.21 ^b	2.38 + 0.46 ^c

* mg Vitamin C equivalent / g fresh weight.

** mg Gallic acid equivalent / g fresh weight.

Means (+ S.D.) with different superscripts in each column are significantly different ($p < 0.05$). Superscript a indicates the group with the highest value and e belongs to the lowest value group

antioxidant capacity analyzed by FRAP assay, followed by *O. sanctum*. In ABTS assay, the plants with the highest antioxidant capacity were *O. basilicum*, followed by *O. sanctum*, respectively. However, results of DPPH assay showed that *O. sanctum*, and then followed by *O. basilicum*, respectively. For overall results, it might be concluded that *O. basilicum* was better sources of antioxidant compounds in this study, followed *O. sanctum*, respectively. In case of total phenolic content analysis, it was found that *O. basilicum*, contained the highest amount of phenolic compounds, followed by *O. sanctum*, respectively.

The correlations between results of antioxidant capacity and TPC analysis are highly significant ($p < 0.01$) as shown in Table 3. These correlations indicate that the phenolic compounds could be the main cause of antioxidant power of *O. basilicum* plant, in accordance with the previous findings that many phenolic compounds in plants are good sources of natural antioxidants (Satoh, 2003 and Singh, 1999). In this study, the contribution of the phenolic compounds content to ABTS values was greater than to FRAP and DPPH values. Thus, ABTS assay may preferably be a better method to study the antioxidant power which is mainly caused by phenolic compounds, than FRAP and DPPH methods, in that order.

Table 3 : Correlation results of three antioxidant capacity assays and total phenolic content (TPC) from all extracts

	Correlation of coefficient	Significance
ABTS	0.825	< 0.01
DPPH	0.973	< 0.01
FRAP	0.683	< 0.01
TPC	1.000	-

Conclusion :

This work shows that *O. basilicum* and *O. sanctum* are good sources of natural antioxidants therefore they should be cultivated in large quantities.

Acknowledgement :

The authors are thankful to Principals of both colleges for their permission.

REFERENCES

AOAC (1997). *Official method of analysis of AOAC international*, 16th Ed., The Association of Analytical Chemists, Arlington.

- Akhtar, M.S. and Munir, M.J. (1989).** Evaluation of the gastric antiulcerogenic effects of *Solanum nigrum*, *Brassica oleracea* and *Ocimum basilicum* in rats. *J. Ethnopharmacol.*, **27**:163-76.
- Arts, M.J.T.J. and Haenen, G.R.M.M.(2004).** Antioxidant capacity of reaction products limits the applicability of the Trolox equivalent antioxidant capacity (TEAC) assay, *Food Chem. Toxicol.*, **42** : 45-49.
- Avesina (1990).** *Law in Medicine*. Interpreter; Sharafkhandy A. Theran: Ministry of Guidance publications; p. 302.
- Brand-William, W. and Cuelier, M. (1995).** Use of free radical method to evaluate antioxidant activity, *Lebensm. Wiss. U. Technol.*, **28** : 25-30.
- Benzie, I.F.F. and Strain, J.J. (1999).** Ferric reducing / antioxidative power assay: direct measure of total antioxidant activity of biological fluids and modified version of simultaneous measurement of antioxidant power and ascorbic acid concentration, *Methods Enzymol.*, **299**:15-27.
- Dasgupta, T., Rao, A.R. and Yadava, P.K. (2004).** Efficacy of basil leaf (*Ocimum basilicum*) on drug metabolizing and antioxidant enzymes, and on carcinogen-induced skin and forestomach papillomagenesis. *Phytomedicine*, **11**:139-151.
- Edris, A.E. and Farrag, E.S. (2003).** Antifungal activity of peppermint and sweet basil essential oils and their major aroma constituents on some plant pathogenic fungi from the vapor phase. *Nahrung*, **47**:117-121.
- Grayer, R.J., Kite, G.C., Goldstone, F.J., Bryan, S.E., Paton, A. and Putievsky, E. (1996).** Intraspecific taxonomy and essential oil chemotypes in sweet basil, *Ocimum basilicum*. *Phytochem.*, **43**:1033-1039.
- Javanmardi, J., Khalighi, A., Kashi, A., Bais, H.P. and Vivanco, J.M.(2002).** Chemical characterization of basil (*Ocimum basilicum* L.) found in local accessions and used in traditional medicines in Iran. *J. agric Food Chem.*, **50** : 5878-5883.
- Jayasinghe, C., Gotoh, N., Aoki, T. and Wada, S. (2003).** Phenolics composition and anti oxidant activity of sweet basil (*Ocimum basilicum* L.). *J. agric Food Chem.*, **51**: 4442-4449.
- Khanna, N. and Bhatia, J. (2003).** Action of *Ocimum sanctum* (Tulsi) in mice: possible mechanism involved. *J. Ethnopharmacol.*, **88**(2-3) : 293-296.
- Lachowicz, K.J., Jones, G.P., Briggs, D.R., Bienvenu, F.E., Wan, J. and Wilcock, A. (1998).** Coventry MJ. The synergistic preservative effects of the essential oils of sweet basil (*Ocimum basilicum* L.) against acid-tolerant food microflora. *Lett. Appl. Microbiol.*, **26** : 209-214.
- Lee, K.G. and Shibamoti, T. (2002).** Determination of antioxidant potential of volatile extracts isolated from various herbs and spices. *J. agric Food Chem.*, **50**:4947-4952.
- Leong, L.P. and Shui, G. (2002).** An investigation of antioxidant capacity of fruits in Singapore markets, *Food Chem.*, **76** : 69-75.
- Mirheidar, H.(1990).** *Herbal Knowledge*. 1st Ed. Islamic Culture Publication; pp. 85-91.
- Opalchenova, G. and Obreshkova, D.(2003).** Comaparative studies on the activity of basil - an essential oil from *Ocimum basilicum* L.-against multi-drug clinical isolates of the genera *Staphylococcus*, *Enterococcus* and *Pseudomonas* by using different test methods. *J Mirobiol. Methods*, **54**:105-110.
- Re, R. et al. (1999).** Antioxidant activity applying an improved ABTS radical cation decolorization assay, *Free Radical Bio. Med.*, **26** :1231-1237.
- Satoh, T. and Sugawara Y. (2003).** Effects on humans elicited by inhaling the fragrance of essential oils: sensory test, multi-channel thermometric study and forehead surface potential wave measurement on basil and peppermint. *Anal Sci.*, **19**:139-146.
- Sen, P. (1993).** Therapeutic potentials of Tulsi : from experience to facts. *Drugs ttggfwdsd News & Views.*, **1**(2): 15-21.
- Singh, S. (1998).** Comparative evaluation of antiinflammatory potential of fixed oil of different species of *Ocimum* and its possible mechanism of action. *Indian J. Exp. Biol.*, **36** :1028-1031.
- Singh, S. (1999).** Evaluation of gastric anti-ulcer activity of fixed oil of *Ocimum basilicum* Linn.and its possible mechanism of action. *Indian J. Exp. Biol.*, **37**:253- 257.
- Singh, S. (1999).** Mechanism of action of antiinflammatory effect of fixed oil of *Ocimum basilicum* Linn. *Indian J. Exp. Biol.*, **37**:248-252.
- Sirkar, N.N.(1989).** Pharmacological basis of Ayurvedic therapeutics. In: *Cultivation and Utilization Of Medicinal Plants*. Editors: Atal CK and Kapoor BM (Published by PID CSIR).
- Suppakul, P., Miltz, J., Sonneveld, K. and Bigger, S.W. (2003).** Antimicrobial properties of basil and its possible application in food packaging. *J. agric. Food Chem.*, **51** : 3197-3207.
- Volak, J. and Plantas Jiri, S. (1997).** *Medicinales*. Interpreter; Zamani S. 3rd Edition. Grund Edition; p. 240.
- Wang, T., Cui, S.Y., Hu, X.L. and Lu, R.H. (2003).** Study on the constituents of volatile oil from *Ocimum basilicum*. *Zhongguo Zhong Yao Za Zhi*, **28** : 740-742.

- Waterhouse, A. (2005).** Folin-Ciocalteu micro method for total phenol in wine. retrieved, from <http://waterhouse.ucdavis.edu/phenol/foolinmicro.htm>.
- Wang, M.F. and Shao, Y. (1998).** Antioxidative phenolic compounds from Sage (*Salvia officinalis*). *J. agric. Food Chem.*, **46**, 4869-4873.
- Yamasaki, K., Nakano, M., Kawahata, T., Mori, H., Otake, T., Ueba, N., Oishi, I., Inami, R., Yamane, M., Nakamura, M., Murata, H. and Nakanishi, T. (1998).** Anti-HIV-1 activity of herbs in Labiatae. *Biol. Pharm. Bull.*, **21** : 829-833.
- Yun, Y.S., Nakajima, Y., Iseda, E., Kunugi, A. (2003).** Determination of anti oxidant activity of herbs by ESR. *Shokuhin Eiseigaku Zasshi*, **44** : 59-62.
- Zargary, A. (1990).** *Medicinal Plants*. 5th Ed. Tehran: Tehran University Publications; pp. 312- 314.

