

Germination improvement through alleviation of seed hardness in *Solanum viarum* Dunal

K. RAMAMOORTHY, R. GEETHA AND K. SUJATHA

Accepted : August, 2009

Key words : Seed hardness, Germination, *Solanum viarum*

Solanum viarum, a member of Solanaceae family, is a hardy, bushy, stout and much branched shrub. The berries of *S. viarum* are rich source of solasodine (Maiti *et al.*, 1964) a phytosteroids used in the pharmaceutical industry, for synthesis of contraceptives, corticosteroids and sex hormones. Some of these steroids also used in the treatment of rheumatoid, arthritis and rheumatoid fever (Kunjithapadam, 1962). *S. viarum* is propagated through seeds and fresh seeds exhibits dormancy due to hard seed coat (Akanda *et al.*, 1996). Hence, studies were initiated to minimize the hard seedness.

Fresh seeds of *S. viarum* were collected from the fully matured fruits and subjected to scarification using commercial sulphuric acid @ 25ml / kg of seed for varying

durations of 5,10,15,20 and 25 min. Seeds were washed four times with water to remove the traces of acid and dried under shade. Scarified seeds were evaluated for germination (ISTA, 1999), root and shoot length, dry matter production and vigour index values (Abdul- Baki and Anderson, 1973). Data were analysed following Snedecor and Cochran (1967).

Fresh seeds without scarification did not germinate. Pingle and Dynansagar (1980) reported dormancy of seeds for an average period of one month and Mullahey *et al.* (1993) recorded enhanced germination in *S. viarum* through mechanical scarification..

In the present study scarified seeds recorded the highest germination percentage (Table 1) when scarified

Table 1 : Effect of scarification in alleviating seed hardness in *Solanum viarum*

Treatments	Germination (%)	Root length (cm)	Shoot length (cm)	Dry matter production	Vigour index
Control	0 (5.73)	0 (0.70)	0 (0.70)	0 (0.70)	0 (0.70)
H ₂ SO ₄ - 5min	45 (42.13)	3.3 (1.82)	3.5 (1.87)	0.012 (0.109)	306 (17.49)
H ₂ SO ₄ - 10min	80 (63.43)	4.3 (2.07)	4.5 (2.12)	0.034 (0.184)	704 (26.53)
H ₂ SO ₄ - 15min	60 (50.76)	3.8 (1.95)	4.1 (2.02)	0.025 (0.158)	474 (21.77)
H ₂ SO ₄ - 20min	50 (45.00)	3.2 (1.78)	3.3 (1.81)	0.012 (0.130)	325 (18.03)
H ₂ SO ₄ - 25min	40 (39.23)	3.0 (2.0)	3.2 (1.78)	0.010 (0.100)	248 (15.74)
S.E. ±	0.57	0.03	0.04	0.004	0.03
C.D. (P=0.05)	1.23**	0.06**	0.08**	0.008**	0.07**

Values in parenthesis are transformed values

** indicates significant of value at P=0.01

Correspondence to:

K. RAMAMOORTHY, Seed Science and Technology Unit, Department of Plant Breeding and Genetics, Agricultural College and Research Institute, MADURAI (T.N.) INDIA

Authors' affiliations:

R. GEETHA AND K. SUJATHA, Seed Science and Technology Unit, Department of Plant Breeding and Genetics, Agricultural College and Research Institute, MADURAI (T.N.) INDIA

with sulphuric acid for 10min (80%) followed by 15min (60%). The root and shoot length (4.3 and 4.5cm, respectively), dry matter production (0.034 mg) and vigour index values (704) were also the highest at 10 min scarification compared to other treatments. Sulphuric acid scarification beyond 15min was injurious and caused reduction in germination and seedling growth.

Hence, scarification with commercial sulphuric acid

@ 25ml/ kg of seeds for 10min. duration is recommended for improving germination of *Solanum viarum*.

REFERENCES

- Abdul Baki, A.A. and Anderson (1973). Vigour determination in soybean seed by multiple criteria. *Crop Sci.*, **13**: 630-633.
- Akanda, R.U., Mullahey, J.J. and Shilling, D.G. (1996). Environmental factors affecting germination of tropical soda apple (*Solanum viarum*). *Weed Sci.*, **44**: 570–574.
- ISTA (1999). International Rules for Seed Testing. *Seed Sci. & Technology (Supplement Rules)*, **27**: 25 – 30.
- Kunjithapadam, S. (1962). Utilization of various raw materials for the production of steroids. In: *Cultivation and Utilization of Medicinal Plants*. Eds. C.K. Atal and B.M. Kapur, Pub. Regional Research Laboratory, Jammu, pp. 51-63.
- Maiti, P.C., Mookerjee, S., Mathew, R and Henery, A.V. (1964). *Solanum khasianum* var *Chaterjeanum* Sengupta, the richest source of Solasodine. *Curr. Sci.*, **33**: 730.
- Mullahey, J.J., Nee, M., Wunderlin, R.P. and Delaney, K.R. (1993). Tropical soda apple (*Solanum viarum*): a new weed threat in subtropical regions. *Weed Technology*, **7**: 783–786.
- Pingle, A.R. and Dynansagar, R. (1980). *Solanum viarum* as a source of solasodine. *Indian Drugs*, **17**: 366- 370.
- Snedecor, G.W. and Cochran, W.G. (1967). *Statistical Methods*. AMEC IOWA State University Press.

