

Alteration of resting period of pollen of five cultivars of apocynaceae by vitamin (vitamin BC) : Further evidence of a criticism of Brewbaker and Kwack (1963), Sudhakaran (1967-Ph. D Thesis) and Saoji and Chitale (1972)*

S.A. SALGARE

Accepted : March, 2010

SUMMARY

Vitamin BC altered the resting period of pollen of 6 series and failed in 4 series of Apocynaceae.

Key words : Palynology, Vitamins, Growth regulators

The phenomenon of the resting period of pollen was reported for the first time by Salgare (1983).

MATERIALS AND METHODS

Pollen of successive flowers (*viz.*, F, F-24, F-48, F-72 series *i.e.* open flowers and the flower buds (which require 24, 48, 72 hours to open, respectively) of 5 cultivars of Apocynaceae *e.g.* red-, pink- and white-flowered cultivars of *Nerium odorum* Soland. and pink- and white-flowered cultivars of *Catharanthus roseus* (L.) G. Don. were collected at the stage of the dehiscence of anthers in the open flowers. Germination of pollen grains of successive flowers was studied by standing-drop technique in the optimum concentrations of sucrose as well as in the optimum concentrations of sucrose supplemented with the optimum concentrations of the vitamin (vitamin BC) (Table 1). The rate of pollen germination of successive flowers was determined by fixing the cultures at one hour intervals. Such preparations were continued for 10 hours. Observations on the germination of pollen were recorded 24 hours after incubation.

RESULTS AND DISCUSSION

Potentiality of pollen germinability was recorded in F series of all the 5 cultivars of Apocynaceae studied. Pollen of F-24 series of red-flowered cultivar of *Nerium odorum* and both the cultivars of *Catharanthus roseus* showed their germination in the optimum concentrations of sucrose. Pollen of F-48 and F-72 series of pink-flowered cultivar of *C. roseus* also showed their

germination in the optimum concentrations of sucrose. Potentiality of pollen germinability in Apocynaceae was observed in 10 out of 20 series investigated (Table 1).

Germination of pollen of F-72 series of pink-flowered cultivar of *C. roseus* *in vitro* culture of sucrose was noted in the present investigation. However, Trisa Palathingal (1990) failed to germinate the pollen of F-72 series of pink-flowered cultivar of *C. roseus* in Brewbaker and Kwack's (1963) culture medium. This proves that the culture medium is also having the bearing on the germination of pollen. This also confirms that Brewbaker and Kwack's (1963) culture medium is not ideal for pollen cultures.

The delay in pollen germination was interpreted by Saoji and Chitale (1972) as being due to the grains not being mature enough to effect pollination, immediately after being shed from the anther. Further they stated that 4-5 hours are required for the complete maturation of pollen grains. It was Salgare (1983) who pointed out for the first time that the pollen require resting period before germination and it was the failure of Saoji and Chitale (1972) who misinterpreted the resting period for pollen maturity. Further he (1983) stated that this resting period differs species to species which is also noted in the present investigation (Table 1). This resting period is altered by different chemicals. Present work as well as the extensive work of Salgare (1983, 86, 2008) made it very clear that Saoji and Chitale's (1972) arguments are superficial and misleading.

Vitamin BC altered the resting period of pollen of 6 series and failed in 4 series of Apocynaceae (Table 1). The vitamin reduced the resting period of pollen of one series, while extended in 5 series.

Sudhakaran (1967) failed to trace out the branched pollen tubes and polysiphonous condition which is fairly common even in diploid pollen grains of *Vinca rosea* L. [*Catharanthus roseus* (L.) G. Don.] which was also

Correspondence to:

S.A. SALGARE, Salgare Research Foundation Pvt. Ltd., Parthamesh Society, Shivaji Chowk, KARJAT (M.S.) INDIA

Table 1 : Effect of vitamin BC on the rate of pollen germination of successive flowers of five cultivars of Apocynaceae

Cultivars	Series	%PV	Conc.			trfpg	
			SC	V	C	T	
<i>Nerium odorum</i>							
Pink-flowered	F	91±0.42	50	10	1	5	
White-flowered	F	61±2.87	50	10	3	5	
Red-flowered	F	61±3.17	20	05	1	5	
Red-flowered	F-24	61±3.17	20	10	1	3	
<i>Catharanthus roseus</i>							
White-flowered	F	89±0.97	20	01	1	1	
White-flowered	F-24	89±0.97	50	10	2	1	
Pink-flowered	F	93. ±0.98	20	10	1	1	
Pink-flowered	F-24	93. ±0.98	50	01	1	6	
Pink-flowered	F-48	93. ±0.98	50	Ng ₂	8	8	
Pink-flowered	F-72	93. ±0.98	80	Ng ₂	Ng ₁	Ng ₁	

C, in control sets time required for germination of pollen in optimum concentrations of sucrose; M, optimum concentrations of vitamin B₂ in mg/ml; Conc, optimum concentrations of sucrose and vitamin B₆; SC, optimum concentrations of sucrose in %; Ng₁, and Ng₂, no germination of pollen even after 10 and 24 hours of sowing respectively; PV, pollen viability; T, time required for germination of pollen in optimum concentrations of sucrose + vitamin BC (in treated sets); trfpg, time required for the germination of pollen in control sets and treated sets in hours.

pointed out earlier by Salgare (1983, 86, 2008).

REFERENCES

- Brewbaker, J.L. and Kwack, B.H. (1963). The essential role of Ca ion in pollen germination and pollen tube growth. *Am. J. Bot.*, **50**: 859-865.
- Salgare, S.A. (1983). Pollen physiology of Angiosperms. Ph.D. Thesis, University, Bombay.
- Salgare, S.A. (1986). Pollen physiology of successive flowers. D.Sc. Thesis, Marathwada University.
- Salgare, S.A. (2008). Alteration of resting period of pollen of five cultivars of Apocynaceae by amino acid (arginine): Further evidence of a criticism of Brewbaker and Kwack (1963), Sudhakaran (1967-Ph.D. Thesis) and Saoji and Chitale (1972) – A Critical Review. *J. Env. Bio-Sci.*, **22**: 323-325.
- Saoji, A.A. and Chitale, S.D. (1972). Palynological studies in *Bauhinia variegata* Linn. *Botanique*, **3**: 7-12.
- Sudhakaran, I.V. (1967). Cytogenetic studies in *Vinca rosea* Linn. Ph.D. Thesis, University, Bombay.
- Trisa, Palathingal (1990). Evaluation of industrial pollution of Bombay by pollen – I. M.Phil. Thesis, University, Mumbai.

