

Effect of different media on the mycelial growth of *Lentinus squarrosulus* Mont.

K. SUBRAMONIAN, S.L. SUBHA AND A. SARAVANA GANTHI

Accepted : September, 2009

SUMMARY

Mushroom constitutes one of the man's natural food for centuries. Now they are used in the production of food, feed, drugs, enzymes, herbicides and insecticides. Among these food and drug are found to be the foremost importance and consumed widely. One of the edible mushroom commonly occur in tropical region is *Lentinus squarrosulus*. The present paper deals with the effect of different media on the mycelial growth of *Lentinus squarrosulus*. Different media like natural media, synthetic media and semi synthetic media were used. Mycelial dry weight was obtained from seven days growth onwards at seven days interval upto 28 days. The result showed that the better mycelial growth was obtained in carrot extract medium among the natural media on 28th day and poor mycelial growth reported in the Carboxy methyl cellulose medium among the semi synthetic media. Among the synthetic media, the basal medium produced better mycelial growth.

Key words : Basidiomata, Gills, Mycelia, Pileus, Stipe

Mushrooms are one of the important bioresources available to the common man especially those who cannot afford to go for protein rich food like mutton, egg, milk etc., Mushrooms are equally supplementing these protein sources, but their availability, rich chemical constituents are not known widely. FAO reported that the average protein consumption per head in developing countries is only half of that of the developed countries. Several attempts have been made to increase the protein production through single cell culture of algae, yeast and many moulds. But these methods are highly expensive, cumbersome and need controlled conditions. In order to bridge the protein gap, mushrooms have come to rescue which has good quality of proteins and major essential amino acids constituting of about 20-45% and also possess amino acids like Lysine, Methionine which are available usually in the animal food. Worldwide there are only dozen species are available for the commercial cultivation. The genus *Lentinus* is commonly occurs in plains of India. Some of the common species are *L. sajor-caju*, *L. cladopus*, *L. crinitus*, *L. prolifer* and *L. squarrosulus*. Among these *L. cladopus* and *L. squarrosulus* are known to be edible. (Corner, 1981; Joly and Perreau, 1977). There is a need to increase the number of edible mushrooms. Information on the nutritive value of tropical

edible mushroom *Lentinus squarrosulus* is scanty. The present study has reported the cellular products of *Lentinus squarrosulus* like total crude proteins, total sugars, reducing sugars, glycogen and lipids content to evaluate the nutritive value of the fungus. For the cultivation of mushroom in submerged culture it is impossible to depend upon a single standard medium. The growth kinetics of *Lentinus squarrosulus* vary in different media. Natarajan and Raman (1983) reported that sawdust-carrot extract medium was the best for the *in-vitro* production of fruiting bodies of *Lentinus edodes*. Inorganic substances such as glass wool, vermiculite and perlite served satisfactorily in supporting production of fruiting bodies in liquid media (Miles and Chang, 1987).

MATERIALS AND METHODS

For the present investigation, fresh healthy *Lentinus squarrosulus* were collected from Tirunelveli hills, Tirunelveli District, Tamil Nadu. The pure culture was maintained on PDA slants at $4 \pm 1^{\circ}$ C in dark and sub-cultured at one month interval. Potato-dextrose agar (PDA), Glucose-Asparagine medium, Carboxy methyl cellulose medium, Sabaraud's medium, Carrot extract medium, Saw dust extract medium (Difco Manual, 1953) Nutrient medium, Basal medium (Srivastava and Bano, 1970), Mushroom synthetic medium (Saha and Samajpati, 1983), Mushroom complete medium (Roy and Samajpati, 1983) were used for the analysis. Triplicates were maintained for each experiment unless otherwise specified.

RESULTS AND DISCUSSION

Different natural media like carrot extract, Saw dust

Correspondence to:

K. SUBRAMONIAN, Department of Plant Biology and Biotechnology, The M.D.T. Hindu College, TIRUNELVELI (T.N.) INDIA

Authors' affiliations:

S.L. SUBHA, Govt. Siddha Medical College, Palayalmkottai, TIRUNELVELI (T.N.) INDIA

A. SARAVANA GANTHI, Department of Botany, Rani Anna Govt. College for Women, TIRUNELVELI (T.N.) INDIA

Table 1 : Mycelial growth* of *Lentinus squarrosulus* on different media

Sr. No.	Types of media	Media	7 th Day*	14 th Day*	21 st Day*	28 th Day*
1.	Natural medium	Carrot extract medium	89.0 (4.5)	319.0 (5.3)	525.0 (6.9)	681.0 (7.0)
2.		Saw dust extract medium	60.0 (4.8)	68.0 (6.3)	90.0 (5.7)	168.0 (6.5)
3.	Semi-synthetic medium	Carboxy methyl cellulose medium	09.0 (6.9)	16.0 (7.2)	24.0 (7.3)	33.0 (7.8)
4.		Nutrient medium	123.0 (5.4)	226.0 (5.8)	277.0 (6.2)	290.0 (6.4)
5.		Potato-dextrose broth	75.0 (5.8)	245.0 (5.20)	349.0 (6.7)	380.0 (7.1)
6.		Sabaraud's medium	126.0 (4.9)	328.0 (7.7)	336.0 (7.5)	408.0 (7.6)
7.		Mushroom complete medium	227.0 (4.9)	436.0 (6.0)	580.0 (4.60)	630.0 (6.7)
8.	Synthetic medium	Glucose asparagine medium	13.0 (6.1)	22.0 (6.0)	33.0 (6.3)	39.0 (6.2)
9.		Mushroom synthetic medium	23.0 (5.1)	77.0 (4.9)	124.0 (5.7)	265.0 (5.8)
10.		Basal medium	44.0 (5.0)	182.0 (5.0)	229.0 (5.0)	360.0 (5.6)

* mg dry weight /flask.

Initial pH 6.0

Incubation period 28 days

Each value mean of triplicate

extract, Semi synthetic media like Carboxy methyl cellulose, Nutrient medium, Potato dextrose broth, Sabaraud's medium, Mushroom complete medium, Synthetic media like glucose-Asparagin medium, Basal medium were used to study the effect of different media on the mycelial growth of *Lentinus squarrosulus*. The initial inoculum was obtained from the fungus growing on PDA plates. Mycelial dry weight was obtained from seven days growth onwards at seven days intervals upto 28 days. The results are presented in Table 1. The results showed that among natural media carrot extract medium had better mycelial growth when compared with Sawdust medium. Among the semi-synthetic media, Mushroom complete medium had the best mycelial growth followed by Sabaraud's medium, Potato dextrose broth and nutrient medium. Compared to the natural and semi-synthetic media, the growth of mycelium in synthetic media was very poor. Among the different media used, Basidiomata initials were produced only in mushroom complete medium

and potato dextrose broth. The change in the pH of different media at different stages of mycelial growth is given in Table 1. In general there was a decrease in the pH on the 7th day and thereafter there was an increase in the pH except in carboxy methyl cellulose medium, where the pH has increased up to 28th day.

The best mycelial growth was obtained in carrot extract medium, whereas abnormal fruiting bodies were produced after 26 days in mushroom complete medium and potato dextrose broth. In a similar study Natarajan and Raman (1980) found the normal fruiting bodies in *Lentinus cladopus* in Sawdust, Carrot extract medium and beetroot extract medium. Abnormal fruiting bodies were produced in most of the other natural and semi-synthetic media. Among the semi-synthetic media the potato dextrose broth was not found suitable for *Lentinus*, but earlier reported suitable for different species of *Pleurotus viz.*, *P. sajor-caju*, *P. florida*, *P. flabellatus* and *P. ostreatus* (Suharban and Nair, 1991).

REFERENCES

- Corner, E.J.H (1981). The Agarics Genera *Letinus*, *Panus* and *Pleurotus* with particular reference to Malaysia species. *Beih Nova. Hedw.*, **68**: 169.
- Difco Manual (1953). *Difco Manual of Dehydrated Culture Media and Reagents for Microbiological and Clinical Laboratory Procedures* (9th Ed.) Difco Laboratories, Detroit.

- Joly, P. and Perreau, J. (1977). Sur quelques Champignons Sauvages consommés au Viet-nam. Travaux-dedies a G.Vietnot-Bourgin: 159-168.
- Miles, G. and Chang, S.T. (1987). Fruiting of *Lentinus edodes* (Shiitake) in liquid media. *Mircen J. Appl. Microbiol. Biotechnol.*, **3** : 103-112.
- Natarajan, K. and Raman, N. (1980). *In vitro* production of fruit bodies in *Lentinus cladopus* in liquid culture. *Indian J. Exp. Biol.*, **18**: 545-547.
- Roy, S.K. and Samajpati, N. (1983). Effects of different media, and incubation periods on the growth of some tropical edible mushrooms. *Indian J. Mycol. Res.*, **21**:45-47.
- Saha, A.K. and Samajpati, N. (1983). Effect of different media and incubation period on the growth of some tropical edible mushrooms. *Indian J. Mycol. Res.*, **21**:49-51.
- Srivastava, H.C. and Bano, Z. (1970). Nutritional requirements for *Pleurotus flabellatus*. *Appl. Microbiol.*, **19**:166-169.
- Suharban, M. and Nair, M.C. (1991). Growth of different species of *Pleurotus* in different media in shake culture. In: Proc. Natl. Symp. on Mushrooms, (ed.) M.C. Nair, Kerala Agriculture University, Thiruvananthapuram, India, pp. 139-140.

