

Effect of *Azospirillum* on fibre quality and yield of irrigated cotton

MUKUNDRAJ B. PATIL

Department of Botany, Late Ramesh Warpudkar A.C.S. college Sonpeth, PARBHANI (M.S.) INDIA
E-mail: - mukundrajbpatil@gmail.com

(Received: June, 2011; Revised: Aug., 2011; Accepted : Sep., 2011)

To study the effect of *Azospirillum* on fibre quality and yield of irrigated cotton, a field experiment was conducted during the year 2003-2004. The field experiments were conducted using Randomized Block Design. Amongst the different strains of *Azospirillum*, Surat strain of *Azospirillum* brought about maximum increment in number of branches and number of bolls, maximum boll weight and highest cotton seed yield. The span length, uniformity ratio, tenacity and elongation percentage (EIG per cent) were increased while the micronaire value and shoot fibre index (SFI) were minimum with the application of Surat strain of *Azospirillum*, which are being considered as average to medium type, good elongation and excellent quality fibres.

Key words : *Azospirillum*, Cotton, Yield, Fiber quality.

B. Patil, Mukundraj. (2011). Effect of *Azospirillum* on fibre quality and yield of irrigated cotton. *Asian J. Bio. Sci.*, 6(2) : 188-190.

INTRODUCTION

Cotton has been one of the main sources of India's economic growth and a foreign exchange earner. Although the country boasts of large area (nearly 9 million hectare), second in yarn production and 3rd in raw cotton production (around 15 million bales annually), the production is far below the world average. Moreover, there has been stagnation in yield level around 300 kg lint/ha, which does not augur in any way any sign of competitiveness (Anonymous, 2002). Seeds inoculated with *Azospirillum* resulted 1.5 times increase in the length of *Flax* fibre (Mikhailouskaya, 2006). Seed yield of *Brassica juncea* (L.) was increased considerably due to application of *Azospirillum* along with neem cake (Khan *et.al.*, 2010). *Azospirillum* not only fixes the nitrogen biologically, but also known to produce growth promoting substances, which favours better growth of crop plants, increase the yield of crop by 5-20 per cent with the saving of nitrogen 40 per cent of the RDF. Taking into account the beneficial role of *Azospirillum*, the present study was undertaken to assess the impact of inoculation on yield and fibre quality of irrigated cotton.

RESEARCH METHODOLOGY

The field experiment was carried out at Research farm, Department of Agronomy, College of Agriculture,

Marathwada Agriculture University, Parbhani. Randomized Block Design was used for the experiment. with three replications. Three most efficient strains (TNAU, HAU and Surat) of *Azospirillum* available in the department of plant pathology were used for inoculation of seed. The cotton variety PH 348 was used for sowing. Sowing was done by dibbling method at row to row 60cm and plant to plant 30cm spacing. Irrigation and standard plant protection were followed. The observations on different yield contributing parameters like number of branches and number of bolls were recorded. For this five plants were selected randomly from each plot. The total number of bolls and branches on each plant were counted physically and average was recorded as number of bolls and number of branches. The seed cotton yield was recorded in each plot after three pickings; yield so recorded in each plot was converted in kg/ha. The representative sample of around 110g of cotton lint each of all treatment and replication was drawn and sent to CICR laboratory, Athwal farm, Surat and data on fibre quality parameters were drawn.

RESULTS AND ANALYSIS

The effect of inoculation of *Azospirillum* on yield and fiber quality parameters of irrigated cotton was studied. The effect of inoculation of *Azospirillum* on

Table 1 : Effect of *Azospirillum* on branches, bolls and seed cotton yield of irrigated cotton

Treatments	No. of branches	No. of bolls	Weight of boll (g)	Seed cotton yield kg/ha
Inoculation of TNAU	24.5	60.80	17.27	1247.5
Inoculation of HAU	26.14	64.23	17.88	1319.7
Inoculation of Surat	27.63	68.36	18.78	1362.8
No Inoculation	19.76	48.90	15.46	1155.7
SE \pm	0.067	0.24	0.055	4.97
CD at 5 %	0.2	0.76	0.16	15.29

Table 2 : Effect of *Azospirillum* on fibre quality parameters of irrigated cotton

Treatments	2.5% Span length	Uniformity ratio	Micro-naire value	Tenacity (g/t) 3.2mm	EIG (%)	Maturity ratio	SFI (W)
Inoculation of TNAU	27.33	50.66	5.02	21.53	6.56	0.83	9.26
Inoculation of HAU	27.3	50.33	5.02	21.52	6.55	0.85	9.53
Inoculation of Surat	27.52	50.91	5.00	21.78	6.66	0.85	8.90
No Inoculation	27.21	49.50	5.09	21.45	6.54	0.85	9.99
SE \pm	0.28	0.26	0.05	0.27	0.02	0.002	0.5
CD at 5 %	NS	NS	NS	NS	NS	NS	NS

NS= Non-significant

branches and bolls was recorded at 90 days after sowing (DAS) and the seed cotton yield was recorded in each plot after three picking. The yield so recorded in each plot was converted in kg/ha. The relevant so obtained are presented in Table 1.

The data (Table 1) clearly indicated that number of branches, weight and number of bolls and seed cotton yield was significantly increased with the inoculation of *Azospirillum*, Surat strain of *Azospirillum* had brought about the maximum number of branches (27.63) and number of bolls (68.36), maximum bolls weight (18.78). It was followed by HAU strain of *Azospirillum*.

The beneficial effects obtained in the present investigation are in full agreement with those reported in the past. Wani *et al.*, (1985) observed consistent increase in yield of pearl millet when inoculated with *Azospirillum lepoferum brasilense*. Anonymous (2002) reported increase in number of bolls, number of branches and 40 per cent increase in seed cotton yield due to application of *Azospirillum* with PSB and PPEM. Martin *et al.*, (2008) recorded increase in wheat yield as a result of increase in the number of spikes and seeds, increase in the kernal weight when inoculated with *Azospirillum brasilense*.

Fibre quality :

The fibre quality of cotton as influenced by inoculation of *Azospirillum* assessed from each plot and the data so obtained are presented in Table 2.

The data on different fiber quality parameters clearly evident that different treatment brought about the changes in fibre quality of cotton. However, these changes could not achieve the level of significance. Of the three strains of *Azospirillum*, Surat strain of *Azospirillum* brought about the maximum span length (27.52), tanacity (21.78g/t) and EIG per cent (6.66 per cent). The fibres obtained due to these treatments are classified as medium type, excellent quality, lowest grade and of good elongation. While the micronaire value (5.00) and shoot fibre index (8.90 W) were minimum with application of Surat strain of *Azospirillum*, which was of superior order and considered as good quality fibre, respectively. Increase in the length of *Flax* fibre due to inoculation with *Azospirillum* was also observed by Mikhailouskaya (2006).

Though, non-significant, still the bioinoculants improved the fibre quality of cotton. The enhancement in fibre quality could not withstand the statistical treatment even though the increase was appreciable. The necessitates in finding out a suitable statistical treatment to such quality parameters. The results obtained in present investigation are in agreement with those reported in the past (Cassman *et al.*, 1990; Phipps *et al.*, 1997).

LITERATURE CITED

- Anonymous (2002).** Annual progress report of technology mission on cotton, mini-mission for the year 2002, pp. 206-219.

- Cassman, K.G., Kerby, T.A., Roberts, B.A. and Brouder, S.M. (1990).** Reassessing potassium requirement of cotton for yield and fibre quality. Proc. of the 1990 Beltwide Cotton Production Conference, pp. 60-64.
- Khan, Irfan, Aquil, Ahmad and Anwar, Masood (2010).** Effect of *Azospirillum* inoculation and organic manure on *Brassica juncea* (L.) *Internet. J. Plant Sci.* **5**(2):669-671.
- Martin, Diaz-Zoritaa Maria, Virginia, Fernandez and Conigiab (2008).** Field performance of aliquid formulation of *Azospirillum brasilense* on dry land wheat productivity. *Eur. J. Biol. Doi.* **10**:1016.
- Mikhailouskaya, N. (2006).** The effect of *Azospirillum brasilense* on *Flax* yield and it's quality. *Plant Soil Environ.* **52**(9):402-406.
- Phipps, B.J., Stevens, W.E., Ward, J.N. and Scales, T.V.(1997).** The influence of mepiquat chloride (PIX) and nitrogen rate upon the maturity and fibre quality of upland cotton. Proc. of the Beltwide Cotton Production Conference, New Orleans, LA, USA. 2, pp. 1471-1472.
- Wani, S.P., Chandrapalah, S. and Dart, P.J. (1985).** Response of pearl-millet cultivars. *Expt. Agri.* **21**:175-182.

**** * ****