

RESEARCH PAPER:

Assessment of replacement cost (on-site cost) of soil erosion in Uva high lands tea plantations of Sri Lanka

■ PRASAD DHARMASENA AND M.S. BHAT

Asian Journal of Environmental Science | December, 2011 | Vol. 6 Issue 2 : 199 -202

Received:

September, 2011

Revised :

November, 2011

Accepted :

November, 2011

SUMMARY

The Uva High lands tea plantations in Sri Lanka represent intermediate zone on agro-climatic classification. This study was conducted to assess the runoff, soil loss and subsequent nutrient losses from Vegetative Propagation (VP) and Old Seedling Tea (OST) plantations of Passara region of Uva high lands in Sri Lanka. Four experimental soil erosion measurement units were installed, 2 each for both land categories of 25 m length and 4 m width during 2010-2011 for one successive year. From these land uses were quantified the following standard methodologies. The annual runoff soil loss of old seedling tea fields were recorded as 25.52 tons/ha/yr and VP fields were calculated as 3.41 tons/ha/yr respectively of Uva regions in Sri Lanka. Loss of N was recorded as 29.34 and 4.80 kg/ha/yr from seedling and VP tea fields, respectively. Loss of P of seedling tea field was observed as 2.10 and P from VP field was 0.92 kg/ha/yr. Loss of K was calculated as 182.4 kg/ha/yr of seedling field and K was assessed in VP as 13.6 kg/ha/yr. Total loss of organic matter was evaluated as 319.01 and 60.03 kg/ha/yr seedling and VP tea fields, respectively. Subsequently, total replacement (onsite) cost of one hectare seedling tea fields was recorded as Rs. 18011.45 and the replacement cost of VP field was Rs.8270.89 with the labour charges for spreading fertilizers and repairing and maintaining costs.

How to cite this paper: Dharmasena, Prasad and Bhat, M.S. (2011). Assessment of replacement cost (on-site cost) of soil erosion in Uva high lands tea plantations of Sri Lanka. *Asian J. Environ. Sci.*, 6(2): 199-202.

Key Words :

Land use,
Seedling, Soil
and nutrient
loss, Runoff,
Uva, NPK,
Organic matter

Land degradations have been a major environmental issue in tea estates of Sri Lanka compared to rubber and coconut plantations. Nearly about 80 per cent of the land is old seedling tea which is often poorly managed (Krishnaraja, 1983). Large tracts of these old seedling tea plantations have been either neglected or left for fallows. It is estimated that about 30 per cent of the entire tea land is marginal or uneconomic in mid country. Long steeps and poor management practices are responsible for severe soil erosion on tea lands (Sivapanal, 1993).

Early plantations industry was under the management of British planters and there were no other parties in the industry with entitlement for the plantations. But tea industry of Sri Lanka today depends on three parties on management systems namely government tea estate, large scale private plantations and small holders. As this research concerned with large scale plantations where can be seen more seedling tea lands where is required as immediate rehabilitation and management. Seedlings tea lands of the large scale

tea plantations (Special reference to the Regional Plantations Companies in Sri Lanka- RPCs) have two types of marginal lands on the definition of Dharmasena (2008). Those are low productive lands, Zero productive lands. Well managed VP tea fields are ecologically and economically stabilized on records maintained by the respective plantations companies.

Study area:

Study sites are based in Passara tea growing region, which is one of controversial regions in environment management of the country and is rested in eastern slopes of the central mountain of the country. It is proved that more abandoned lands in Badulla tea region district are observed in Passara region compared with other tea growing areas of the district where boarded to two dry zone extremes.

EXPERIMENTAL METHODOLOGY

Four soil sedimentation plots were

Author for

Correspondence -

**PRASAD
DHARMASENA**

P.G. Department of
Geography and Regional
Development,
University of Kashmir,
SRINAGAR
(J. & K.) INDIA

See end of the paper
for **Coopted authors**

established in Passara of Sri Lanka in order to measure the physical soil loss of two land use categories namely, VP and seedling tea lands for the period of September 2010 - September 2011. Two soil sedimentation plots were established in Dammeria a estate of Passara region and another two were established in Adawatte estate of Passara region. Average elevation of experimental block in Passara region is between 996 m and 1120 m. Soil of the site is Red Yellow Podzolic (RYP) in the experimental sites. The general elevation was laid between 889 m and 1098 m of these experimental sites. Slope gradient of both experimental sites were 20-30 per cent. More of mid country tea estates are placed between 20 per cent and 30 per cent land slope category (Central Environment Authority of Sri Lanka, 1998). Besides, steep slope lands are not allowed to be used for any development activity by the Central Environmental Authority and Tea Research Institute of Sri Lanka.

All these four experimental plots were designed along the line of the experiment carried out by Jinza (1981). He designed soil erosion sedimentation plots varying for 60m² to 200m². Even though, this particular experiment was based on sedimentation plots introduced by Jinza(1981), Size of sedimentation plot was standardized for 100 m² of 25 m length and 4 m width in the area with a regular shape and sedimentation pit was 3 m x 1.5 m x 2 m. Entire sedimentation plot was completely covered by using high gauge galvanize sheets for ensuring the prevention of over flowing run off or water going out from the plot. Runoff water from the sedimentation plot was collected in the sedimentation pits which have been well covered using white durable polythene.

The sediments were dry after collecting of soil, if collecting period was rainy; until, it comes to normal moisture percentage and method is to weigh the sediment in the pit by using a commercial digital scale with acceptable accuracy. Data also were collected once in two months. Deduct the moisture content to get the exact dry weight of moisture-free soil collected per pit, the results in the soil lost from 100 m² of area and is converted to determine the quantity lost from one hectare. Samples collected from each plot were tested in the laboratory to determine total loss of N, P, K and SOM etc. Fig. 1 is an example of the experiment plots installed in Dammeria A estate of Passara tea region. Precipitation is measured by standard rain gauges of 200 mm tunnel diameter and recording gauges. The total amount of precipitation and the time of the start and end of each rainfall season were recorded. The distribution of precipitation with respect to time was observed mainly by recording gauges, and in places without such experimental facilities the time-dependency is obtained by taking manual readings at certain intervals.



Fig. 1: Sedimentation pits which have been well covered using white durable polythene

EXPERIMENTAL FINDINGS AND DISCUSSION

Soil loss measurement plots systems used to measure soil loss in Passara region for selected land used types. Normal agricultural practices have been carried out during the experimental period. But applications of manure were closed before 6 months until experiments were completed, as it is required to ensure annual loss of NPK and organic carbon. The function of soil loss measurement unit was started in same date of the region.

All land use types could be selected in one division for installing soil loss measurement units as it was required to verify the degree of issue in under the same independent variables. Table 1 elaborates the soil loss of 100m² land area for the tested period.

Soil nutrients loss is the major soil degradation issue in mountaineous plantations. This hazard is occurred due to soil degradation through soil erosion. Soil nutrient depletion is a major concern in terms of resource utilization and optimization of production in the tea plantations. Soil and nutrient losses from plantation lands are responsible for pollution of surface waters and this reduces the productive capacity of land in direct and formation of unfavourable ecosystem is another issue related to plantation lands.

Parallel to soil loss measurements test carried out in two estates of the Uva region, has been tested nutrients loss of different land uses systems. During the period of

Table 1 : Soil loss at 100m² territory in Uva region for the year 2010-2011

Region	Plot No.	Land use type	Duration	Rainfall(mm)	Soil loss (kg)
Passara	1	seedling tea	Sep-Nov	394.11	47.7
Passara	1	seedling tea	Nov-Jan	562.23	35.87
Passara	1	seedling tea	Jan-Mar	184.54	32.2
Passara	1	seedling tea	Mar-May	587.5	77.38
Passara	1	seedling tea	May-Jul	165.2	27.3
Passara	1	seedling tea	Jul--Sep	138.33	35.07
Total kg for 100m ²					255.52
Total per hectare					25.52 tons/yr /ha
Region	Plot No.	Land use type	Duration	Rainfall(mm)	Soil loss (kg)
Passara	2	VP	Sep-Nov	394.11	5.4
Passara	2	VP	Nov-Jan	562.23	8.67
Passara	4	VP	Jan-Mar	184.54	5.46
Passara	4	VP	Mar-May	587.5	7.34
Passara	4	VP	May-Jul	165.2	4.06
Passara	4	VP	Jul--Sep	138.33	3.21
Total kg for 100m ²					34.14
Total per hectare					3.414 tons/yr/ha

2010 -2011, soil loss was measured one time for a two months period and it came total of six bulks for determination of total eroded extent during the tested period. Well mixed two representative samples were collected from the soil eroded pits and 12 samples were total. After testing, these 12 samples to determine total nutrients loss, it was calculated average NPK percentage, pH, texture and OM percentage during the period. Table 2 explains the nutrients content of silted soils of the pits of two lands uses of the region.

Table 2 shows the average nutrients loss of different land use types. Soil tests were executed at six times in different 12 samples. Eventually average nutrients loss of 12 samples was calculated for representing annual loss of nutrients and finally current market prices were multiplied by total nutrients loss to determine the total annual onsite cost of VP and seedling tea fields in Uva region of Sri

Table 2 : Nutrients percentage of silted soil in measurement pits

Region	Composition					
	N (%)	P (ppm)	K (ppm)	OM (%)	Texture	pH
Passara						
Seedling field	0.115	17	45	1.25	SC	4.5
VP Field	0.141	36	96	1.76	SC	4.6

Lanka (Table 3). Table 3 illustrates the total nutrients loss in Sri Lankan rupees.

Conclusion and recommendation:

These figures illustrate that onsite cost and nutrients loss were very high in land use of seedling type tea

Table 3 : Total nutrients loss and current nutrients prices (yr/ha)

Items	VP	Seedling	Price per kg(Rs)
N(kg)	4.8	29.34	35
P(kg)	0.92	2.1	18
K(kg)	13.6	182.4	31
OM	60.03	319.01	10

Table 4 : Total onsite cost (replacement cost) for VP and seedlings tea plantations (yr/ha)

Items	VP	Seedling
N(Rs.)	168	1026.9
P(Rs.)	16.56	37.8
K(Rs.)	421.6	5654.4
OM(Rs.)	600.8	3190.1
Labor cost(Rs.)	2500	2500
R and M(Rs.)	4564	5602.25
Total (Rs.)	8271.89	18011.45

plantations. Approximate seedling tea extent of the region was reported as 11000 in hectares (Statistics division, 2009). Total onsite cost of seedling tea fields of the region was as high as Rs: 198125950.00.

Vegetation cover is the ideal system for controlling the issue related tea garden of Uva region of Sri Lanka. It will also help to maintain good tea ecosystems of the region. Hence, practicing of tree based mix cropping models in this region should be practiced as remedy for the issue in order to maintain the stability and sustainability of tea eco-system. Relevant authorities should take immediate action to re-introduce biological soil conservation systems

which have been practiced before privatization of the industry in Sri Lanka.

Acknowledgement:

Authors are grateful to Mr. Jayampathy Molligoda, Director of Bogawantalawa Plantations Ltd. and Mr. Nilan Liyanage, Senior Superintendent of Telbedde Estate of Balangoda Plantations Ltd. of Sri Lanka for their valuable contribution to the study. Authors are also thanking to all officials of Finlays' Plantations PLC of Sri Lanka.

COOPTED AUTHORS-

M.S. BHAT, P.G. Department of Geography and Regional Development, University of Kashmir, SRINAGAR (J.&K.) INDIA

REFERENCES

Central Environment Authority of Sri Lanka. (1998). *Optimal land use in hill country of Sri Lanka*. Colombo: CEA.

Dharmasena, P. (2008). *Agroforestry models to improve economic viability of marginal tea lands*.

Krishnaraja, P. (1983). Soil loss and conservation planning in tea plantations in Sri Lanka. *Natural System for Development*, pp.141-161.

Jinza, MOU (1981). The establishment of experimental plots for studying runoff and soil loss in the rolling loess regions of China. *Erosion and Sediment Transport Measurement (Proceedings of the Florence)* (pp. 467-477). Zhengzhou,: IAHS Publishers.

Sivapanal.P. (1993). Tea industry in Sri Lanka. International Symposium on tea science and human health (pp. 77-85). Culculla-India: Tea Research Association.

Statistics Division. (2009). *Tea statistics*. Colombo: Sri Lanka Tea Board .

