

RESEARCH NOTE

Screening of pea varieties, germplasm lines and genotypes against pea wilt (*Fusarium oxysporum* f.sp. *pisi*)

■ D.P. KULDHAR, S.IL. BADGUJAR AND UTPAL DEY*

Department of Plant Pathology, College of Agriculture, Marathwada Krishi Vidyapeeth, PARBHANI (M.S.) INDIA

ARTICLE INFO

Received : 12.07.2012

Accepted : 27.02.2013

Key Words :

Pea wilt,

Screening,

F. oxysporum

f.sp. *pisi*

*Corresponding author:
utpaldey86@gmail.com

ABSTRACT

Field experiment has been carried during *Rabi*, 2009, to find out pea varieties, germplasm lines and genotypes against pea wilt incited by *Fusarium oxysporum* f.sp. *pisi*. Among the pea varieties tested, Arkel was found most susceptible with significantly highest mean wilt incidence (22.66%), and this was followed by Latur local-1 (29 %) and Latur Local-2 (28%). Soldier was found moderately susceptible with mean wilt incidence of 19 per cent. Thus, pea variety Soldier may be preferred for sowing during first fortnight of October, so as to minimize the yield losses due to wilt incidence.

How to view point the article : Kuldhar, D.P., Badgular, S.L. and Dey, Utpal (2013). Screening of pea varieties, germplasm lines and genotypes against pea wilt (*Fusarium oxysporum* f.sp. *pisi*). *Internat. J. Plant Protec.*, 6(1) : 209-210.

Pea (*Pisum sativum* L.) is one of the important pulse crops grown in Maharashtra. Peas are cultivated for the fresh green seeds, tender green pods, dried seeds and foliage and cooked as a vegetable, marketed fresh, canned and frozen. Pea is affected by several plant pathogens including fungi, bacteria and viruses that cause serious diseases. Among the fungal diseases, wilt incited by *Fusarium oxysporum* is considered as one of the most devastating diseases of the pea, which results in heavy yield losses.

Wilt (*F. oxysporum*) is one of the most important diseases and reported to occur worldwide in the countries viz., India, Bangladesh, Brazil, Philippines, South Australia, Taiwan, Thailand, Tropical Africa, France, USA, Pakistan, China, Russia, Canada and many other countries. In India, wilt (*F. oxysporum*) was reported to occur and cause heavy quantitative and qualitative losses in pea, chickpea, pigeonpea, lentil and many other important pulse crops grown in the state of Maharashtra, Karnataka, Andhra Pradesh, Orissa, Tamil Nadu and Madhya Pradesh (Raabe *et al.*, 1981 and Grewal, 1988).

Pea cultivation throughout the country in general and in the region of Marathwada and Western Maharashtra particularly has been facing the serious menaces of wilt

incidence. Most of the pea cultivars under cultivation are highly susceptible to the wilt. Hence, it was necessary to find out the disease resistant/tolerant sources for successful management of the disease.

Keeping in view, the economic importance of the crop and yield losses caused by *Fusarium* wilt in pea, present study was undertaken.

To identify the sources of wilt resistance in the varieties, cultivars and germplasm lines of pea, screening was undertaken in screen house at the Department of Plant Pathology during *Rabi*, 2009.

Screening of available pea varieties, germplasm lines, genotypes was conducted in *F. oxysporum* f. sp. *pisi* sick soil in the pot by adding the inoculum. A total of 12 genotypes were sown in pot (10 seed/ pot). The cultivars, varieties and germplasm lines of pea evaluated were Soldier, Arkel, Latur local-1, KS-205, Latur local-2, KS-210, VP-215, Arka Ajit, PMR-53, IP-3, VP-433, and VP-434.

Observations on number of plants wilted from each genotype were recorded at 30, 45, 60, 75 and 90 days after sowing. The per cent wilt incidence was calculated on the basis of initial plant count and total number of wilted plants in each genotype and graded as follows:

Table 1: Reaction of pea germplasm lines against *F. oxysporum* f. sp. *pisi*

Sr. No.	Genotype	Mean per cent disease incidence	Disease reaction
1.	Soldier	19.00	MR
2.	Arkel	32.66	S
3.	Latur local-1 (small seeded)	29.00	MS
4.	KS- 205	17.00	MR
5.	Latur local-2 (large seeded)	28.00	MS
6.	KS- 210	15.00	MR
7.	VP- 215	18.00	MR
8.	Arka Ajit	27.33	MS
9.	PMR- 53	19.00	MR
10.	IP- 3	48.10	S
11.	VP-433	19.00	MR
12.	VP-434	20.00	MR

R : Resistant, MR : Moderately resistant, MS : Moderately susceptible, S : Susceptible, HS : Highly susceptible

Reaction	Per cent wilting (mortality)
– Resistant (R)	0 – 10 % mortality
– Moderately resistant (MR)	10.1 -20 % mortality
– Moderately susceptible (MS)	20.1 -30 % mortality
– Susceptible (S)	30.1 -50 % mortality
– Highly susceptible (HS)	Above 50 % mortality

A total of 12 pea germplasm lines/varieties/cultivars were screened by following the sick plot technique developed by Nene *et al.* (1981), which was proved to be the most efficient and quick. On the basis of per cent disease incidence, germplasm lines, cultivars were categorized as resistant, moderately resistant, moderately susceptible, susceptible and highly susceptible by using the disease rating scale. The results obtained on per cent disease incidence of different germplasm lines evaluated against *F. oxysporum* f. sp. *pisi* on sick pot developed in screen house are presented in Table 1.

The results of Table 1 indicate that, among 12 germplasm lines/genotypes and varieties of pea, none was found resistant or tolerant to the disease. The mean disease incidence ranged from 15.00 to 48.10 per cent.

Of the 12 germplasm lines/genotypes of pea screened under screen house condition seven lines *viz.*, Soldier, KS-205, KS-210, VP-215, PMR-53, VP-433 and VP-434 were found moderately resistant with disease incidence in the range of 15 to 20 per cent. Three lines *viz.*, Latur local-1 (small seeded), Latur local-2 (large seeded) and Arka Ajit were found moderately susceptible with disease incidence in the range of 27.33 to 29.00 per cent and two line *viz.*, Arkel and IP-3 were found susceptible with the disease incidence of about 32.66 and 48.10 per cent, respectively.

Thus, majority of the pea cultivars/varieties, genotypes

screened and presently under study were found moderately resistant or moderately susceptible to the wilt disease. Very scanty literature is available on pea germplasm screening against *F. oxysporum* f. sp. *pisi*. However, Utikar (1965), Crompton and Gaulten (1974), Sen and Mujumdar (1974) and Verma and Dohroo (2002) had attempted the pea germplasm screening against *F. oxysporum* f. sp. *pisi*.

REFERENCES

- Crompton, M.S. and Goulden, D.S. (1974).** New peas resistant to wilt. *New Zealand. J. Agric.*, **128** (1): 12-13.
- Grewal, J.S. (1988).** Diseases of pulse crops: An overview. *Indian Phytopath.*, **41** (4): 1-14.
- Nene, Y.L. and Kannaiyan, J. and Reddy, M. V. (1981).** Pigeonpea diseases: resistance-screening techniques. ICRIASAT. Information Bulletin No. 9. Documentation. International Crops Research Institute for the Semi-Arid Tropics, Patancheru (A.P.) INDIA.
- Raabe, R.D., Connors, I.L. and Martinez A.P. (1981).** Checklist of plant diseases in Hawaii: including records of microorganisms, principally fungi, found in the state. Hawaii Institute of Tropical Agriculture and Human resources (CTAHR), Information Text Series 022.: 313.
- Sen, Bineeta and Majumdar, M. (1974).** Resistance to Fusarium wilt in garden peas. *Indian Phytopath.*, **27**:70-71.
- Utikar, P.G. (1965).** Further studies on Fusarium wilt of peas (*Pisum sativum*. L.) in Maharashtra State. M.Sc. (Ag.) Thesis, Poona University. Poona, M.S. (INDIA).
- Verma, S. and Dohroo, N.P. (2002).** Studies Fusarium wilt of pea (*Pisum sativum* L.) *Thesis Abstract*, **28** (2) : 132-133.
