

Effect of use of polythene mulch on yield of groundnut as tested in the fields of farmer

N.B. MALEKAR AND P.S. ATAWARE

Received : November, 2010; Revised : January, 2011; Accepted : April, 2011

SUMMARY

Groundnut is an important oilseed crop of the country. The yield of groundnut though varies from variety to variety, the cultivation practices also affect much in case of groundnut. One of the major practice which is supposed to increase yield of groundnut is use of Polythene mulch on raised beds especially in summer groundnut. The polythene mulch technology was tested in summer groundnut in the fields of farmers so as to see the effect on yield of the crop. 7 micron transparent polythene sheet was used on raised beds of 75 cm top. Different plots belonging to different farmers at different locations irrespective of the variety and other package of practices they are using, were selected for study to avoid environmental and experimental error. Each farmer's plot was divided into control and experimental plot. The cultural practices except raised beds and polythene mulch were same as that of control in each plot. Observations were recorded for yield. Mean yield level of the control plots was 19.84 q/ha while that of experimental was 23.48 q/ha. There was significant rise in the levels of yield from approximately 10 to 25 % (average 18%) after using polythene mulch for groundnut crop, depending upon individual difference in other practices and management of different farmers.

Malekar, N.B. and Atakare, P.S. (2011). Effect of use of polythene mulch on yield of groundnut as tested in the fields of farmer. *Internat. J. Plant Sci.*, 6 (2): 267-269.

Key words : Groundnut yield, Polythene mulch, Raised beds

Groundnut is one of the major oil crop of India. Oil and oilseed prices are reaching high and if the yield per hectare increases, surely farmer will get benefited and the oil availability will be more. The yield of groundnut though varies from variety to variety, the cultivation practices also affect much in case of groundnut. One of the major practice which is supposed to increase yield of groundnut is use of polythene mulch on raised beds. The polythene mulch technology was tested so as to see the effect on yield of the crop.

MATERIALS AND METHODS

7 micron transparent polythene sheet with one inch diameter holes at 20 cm by 20 cm distance was used for study for treatment. The breadth of the paper was 100 cm. Raised beds of 75 cm top and 30 cm wide furrow in-between were used in both control and treatment plots. Total 4 lines were sown on one raised bed.

Correspondence to:

N.B. MALEKAR, Department of Agricultural Botany,
College of Agriculture, Bahubali, KOLHAPUR (M.S.)
Email : nbmalekar@yahoo.in

Authors' affiliations:

P.A. ATAWARE, School of Agricultural Sciences,
Yashwantrao Chavan Maharashtra Open University,
NASHIK (M.S.) INDIA

Locale of the study:

Different plots belonging to different farmers at different locations irrespective of the variety and other package of practices they are using, were selected for study to avoid environmental and experimental error.

The farmers were convinced to use this new technique in their plots. The field was divided in two parts as, in first part all the practices were commonly used by the farmer(control plot) where as in second raised beds and polythene mulch technology was used.

The fertilizer doses and other practices were kept common in both control and experimental plots. Maximum farmers offered less area for this experiment and major portion for the regular package of practice.

Control plot:

Each plot with raised bed was divided into two parts. Beds in one of the part were not covered with polythene mulch. This plot was considered as control

Treatment plot:

Beds in second *i.e.* remaining part of the plot were covered with 7 micron polythene paper. This plot was considered as treatment plot

Cultivation practices:

Except polythene mulch, all other practices were kept common in both control and treatment plot

No. of plots selected:

Total 44 plots were selected

Record of observations:

The observations were recorded for yield of control plot and treatment plot.

Statistical analysis:

Z test was applied to test the data for its significance

RESULTS AND DISCUSSION

The dry weight yield of the pods was recorded for control plots and experimental plots of the farmers. As the whole experiment was carried out at farmers field, the plot sizes are different for each farmer. Also, many farmers have preferred to allot less area to the experimentation. For making the data homogeneous, yield of the control plot and experimental plot was converted to yield per hectare (Table 1).

The data for yield in control and treatment plot were analyzed using z test. (Table 2) The test was found significant as Z calculated value (8.965553115) is more than Z table value (1.96 at 0.05 alpha value).

This clearly indicates that because of use of polythene mulch for The mean for control plot was 19.84 while of treatment plot was 23.48. The increase in the yield of the plot covered with polythene mulch was approximately 18 per cent.

The technology of use of polythene mulch was introduced by China. Because of its contribution in increasing groundnut production in China, polythene mulching is called as the "White Revolution". Polythene mulching was introduced in China in 1978 from Japan. In China, the polythene mulching was found to increase yield between 18 to 49% depending on soil fertility conditions.

Golombek *et al.* (1995) in their article "Effect of transparent polythene mulch during the reproductive stages of groundnut." published in International Arachis Newsletter of ICRISAT said that "Applying mulch from 50-113 d after sowing increased the mature seed DW/plant by 24%, mainly due to an increased number of seeds/plant."

Maurya and Lal (1981), in their article "Effects of different mulch materials on soil properties and on the root growth and yield of maize (*Zea mays*) and cowpea (*Vigna unguiculata*)" said that "translucent polythene and straw mulch materials yield more than unmulched

[*Internat. J. Plant Sci.*, 6 (2); (July, 2011)]

Table 1 : Observations recorded for yield of groundnut						
Observations Recorded:						
Sr. No.	Control			Experimental		
	Plot size (ha)	Yield (q)	Yield (q/ ha)	Plot size (ha)	Yield (q)	Yield (q/ ha)
1.	0.4	8.44	16.66	0.2	4.8	24
2.	0.2	3.56	17.8	0.2	4.37	21.85
3.	0.4	8.06	20.15	0.2	5.06	25.3
4.	0.4	9	22.5	0.4	10.44	26.1
5.	0.1	2.4	24	0.1	2.75	27.5
6.	0.3	5.25	17.33	0.2	4.04	20.28
7.	0.15	2.5	16.66	0.15	3.15	21
8.	0.3	5.5	18.33	0.1	2.05	20.5
9.	0.4	8.2	20.5	0.2	4.81	24.05
10.	0.2	4.3	21.5	0.1	2.59	25.9
11.	0.2	3.64	18.2	0.2	4.26	21.3
12.	0.25	4.15	16.6	0.15	2.77	18.46
13.	0.2	4	20	0.2	4.56	22.8
14.	0.8	16.74	20.92	0.2	4.82	24.1
15.	0.5	11.54	23.08	0.1	2.67	26.7
16.	0.4	7.25	18.12	0.4	8.88	22.2
17.	0.6	9.5	15.83	0.1	1.86	18.6
18.	0.2	360	18	0.1	2.12	21.2
19.	0.4	7.55	18.87	0.6	13.58	22.63
20.	0.1	1.85	18.5	0.3	6.06	20.2
21.	0.5	11.1	22.2	0.1	2.48	24.8
22.	0.3	5.12	17.06	0.1	1.96	19.6
23.	0.2	3.05	15.25	0.2	3.75	18.75
24.	0.2	5.6	28	0.2	7.4	37
25.	0.4	8.12	20.3	0.4	9.42	23.55
26.	0.3	5.85	19.5	0.1	2.39	23.9
27.	0.3	6.73	22.43	0.1	2.63	26.3
28.	0.2	3.82	19.1	0.1	2.38	23.8
29.	0.4	8.18	20.45	0.1	2.45	24.5
30.	0.4	12.17	30.42	0.2	6.98	34.9
31.	0.3	690	23	0.1	2.69	26.9
32.	0.7	12.62	18.02	0.1	2.27	22.7
33.	1.1	18.15	16.5	0.3	6.12	20.4
34.	0.25	4.48	17.92	0.25	5.22	20.88
35.	0.2	3.88	19.4	0.2	4.66	23.3
36.	0.2	3.77	18.85	0.4	8	20
37.	0.4	8.86	22.15	0.4	9.88	24.79
38.	0.35	5.9	16.85	0.2	3.84	19.2
39.	0.3	4.9	16.33	0.1	1.87	18.7
40.	0.2	4.81	24.05	0.2	5.56	27.8
41.	0.3	6.05	20.16	0.2	4.83	24.15
42.	0.1	1.96	19.6	0.3	6.96	23.2
43.	0.4	9.12	22.8	0.4	10.68	26.7
44.	0.4	7.63	19.07	0.2	4.55	22.75
		Total	872.96			1033.24
		Mean	19.84			23.48

Table 2 : Statistical analysis of the data

	z-Test	
	Control plots	Treatment plots
Mean	19.94090909	23.48272727
Known variance	3.071592889	3.795153803
Observations	44	44
z calculated = $ - 8.965553115 = 8.965553115$		
z table value at 5% =	1.96	

To be significant at an alpha of .05, the z-score must exceed 1.96

ridges and transparent mulch”

Also, there are farmers in Maharashtra (Shri Jaykumar Gunde, Pattankodoli, Dist. Kolhapur) who are using this technique since last 17 years (from 1993) and had got 20 to 50 per cent increase in the yield of summer groundnut, depending upon other situations and package of practices.

Thus the technology though not commonly used, is easily applicable and seems to be beneficial so as to increase groundnut yield especially in summer groundnut

REFERENCES

- Ghosh, P.K., Devi Dayal, Bandyopadhyay, K.K. and Mohanty, M. (2006) Evaluation of straw and polythene mulch for enhancing productivity of irrigated summer groundnut. *Field Crops Res.*, **29** (2-3) : 76-86
- Maurya, P.R., and Lal, R. (1981). Effects of different mulch materials on soil properties and on the root growth and yield of maize (*Zea mays*) and cowpea (*Vigna unguiculata*) Original Research Article. *Field Crops Res.*, **4**: 33-45.
- Golombek, S.D., Talwar, H.S. and Johansen, C. (1995). Effect of transparent polythene mulch during the reproductive stages of groundnut. *Internat. Arachis Newsletter*, **15** : 68-69.

