

Effect of age of seedlings under different system of rice intensification (SRI)

B.N. MANJUNATHA*, R. BASAVARAJAPPA¹, SIDDARAM, A.S. POLICEPATIL,
H. YOGESHAPPA² AND K.N. KALYANAMURTHY

Department of Agronomy, College of Agriculture, University of Agricultural Sciences, G.K.V.K., BANGALORE
(KARNATAKA.) INDIA

ABSTRACT

A field experiment was conducted at Agricultural Research Station, Siruguppa, Karnataka, during the *Kharif*, 2005 to study the influence of age of seedlings under different system of rice intensification. Modified SRI method resulted in significantly higher grain yield (6342 kg ha⁻¹) when compared to other methods. Younger seedlings of 9 days (6071 kg ha⁻¹) and 12 days (6018 kg ha⁻¹) produced significantly higher grain yield than the other aged seedlings viz., 15 days (5792 kg ha⁻¹), 18 days (5771 kg ha⁻¹) and 21 days (5781 kg ha⁻¹). The interaction effects between methods of planting and age of seedlings were found to be non significant.

Key words : System of rice intensification, Grain yield, Age of seedlings

INTRODUCTION

To assure food security in the rice-consuming countries of the world, those countries will have to produce 50% more rice with improved quality to meet consumers' demand by 2025. This additional rice will have to be produced on less land with less water, less labour, and fewer chemicals. Rice (*Oryza sativa* L.) is an important crop of Tungabhadra project area and is cultivated over an area of 0.1 m ha. Timely planting and use of appropriate aged seedlings for transplanting are important non cash inputs for realizing the higher productivity in rice (Pattar *et al.*, 2001).

Transplanting younger seedlings, preferably 8-15 days old before the plants enter their fourth phyllochron of growth, planting the seedlings singly rather than in clumps of 3-6 plants, and keeping the paddy soil moist but not continuously saturated during the plants' vegetative growth phase. Transplanting of less than 15 days old seedlings was found better. Further farmers are very often to use 35 to 50 days old seedlings in place of recommended 25 to 30 days old seedlings and such information under TBP situations is not available (Balasubramanian *et al.*, 1977). It is important to select suitable age rice seedlings contributing towards increased rice production and resolve the problem posed by higher temperature and water depth inflicting seedling mortality. This has resulted in low population per unit area just after transplanting as well as causing huge economic losses. Hence, the present investigation was carried out to study the influence of age of seedlings under different system of rice intensification.

MATERIALS AND METHODS

A field experiment was carried out during the *Kharif*, 2005 at Agricultural Research Station Siruguppa, Karnataka. The soil was deep black clay in texture, neutral in pH (8.22) and low in electrical conductivity (0.28 dS/m). It was low in available nitrogen (265 kg ha⁻¹), high in available phosphorus (30.5 kg ha⁻¹) and medium in available potassium content (365 kg ha⁻¹). There were 15 treatment combinations comprising of three methods of planting (M₁- Normal method, M₂- Recommended SRI method, M₃- Modified SRI method) as main treatments and five age of seedlings (9, 12, 15, 18 and 21 days) as sub treatments and were laid out in split plot design with three replications. The gross plot size was 4 m x 3 m. IET-16933 was used as test variety. The spacing followed was 20 cm x 10 cm (M₁), 25 cm x 25 cm (M₂ and M₃). The crop received a fertilizer dose of 150:75:75 kg NPK ha⁻¹. Full dose of P₂O₅ and K₂O and 50 per cent of nitrogen was applied at the time of transplanting and remaining 50 per cent of nitrogen was applied in two equal splits at 30 and 55 days after transplanting. All the recommended cultivation practices were followed. The crop was harvested after attaining physiological maturity.

RESULTS AND DISCUSSION

The results obtained from the present investigation as well as relevant discussion have been presented under following heads:

Methods of planting :

Methods of planting had significant influence on the

* Author for correspondence.

¹Department of Agronomy, University of Agricultural Sciences, DHARWAD (KARNATAKA)

²Department of Soil Science and Agricultural Chemistry, College of Agriculture, University of Agricultural Sciences, G.K.V.K., BANGALORE (KARNATAKA)

Table 1 : Total number of panicles, panicle length, number of grains per panicle and test weight /1000 grain weight as influenced by methods of planting and age of seedlings

	Total number of panicles per hill					Panicle length (cm)					Number of grains per panicle					Test weight /1000 grain weight (g)								
	S ₁	S ₂	S ₃	S ₄	S ₅	Mean	S ₁	S ₂	S ₃	S ₄	S ₅	Mean	S ₁	S ₂	S ₃	S ₄	S ₅	Mean	S ₁	S ₂	S ₃	S ₄	S ₅	Mean
M ₁	19.32	19.10	17.54	17.30	17.67	18.19	19.50	19.73	18.77	18.20	18.57	18.95	129.60	129.53	121.27	120.53	122.40	124.67	19.97	19.80	19.20	19.30	19.20	19.49
M ₂	63.12	62.30	59.71	59.58	59.05	60.75	21.87	22.10	21.20	22.13	20.83	21.63	172.53	173.67	155.27	158.73	158.30	163.70	21.00	20.83	20.37	20.27	20.10	20.51
M ₃	64.35	63.12	60.12	60.23	59.80	61.52	21.83	21.90	20.90	20.37	20.83	21.17	175.87	171.20	167.60	161.33	160.27	167.25	21.27	21.10	20.63	20.67	20.50	20.83
Mean	48.93	48.17	45.79	45.70	45.50	47.50	21.07	21.24	20.29	20.23	20.08	21.17	159.33	158.13	148.04	146.87	146.99	167.25	20.74	20.58	20.07	20.08	19.93	
	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)
Main treatments (M)	0.56	2.22	0.31	0.31	0.21	0.21	0.87	3.44	0.09	0.36														
Sub treatments (s)	0.81	2.38	0.23	0.23	0.68	0.68	1.31	3.83	0.05	0.14														
S at same M	1.41	NS	0.40	0.40	NS	NS	2.27	NS	0.08	NS														
M at same or different S	1.38	NS	0.47	0.47	NS	NS	2.21	NS	0.12	NS														

M₁ – Normal method (Application of RDF through inorganics with FYM @ 10 t ha⁻¹ and transplanting at 20 cm x 10 cm spacing)
M₂ – Recommended SRI (Application of RDN through organics and transplanting at 25 cm x 25 cm spacing)
M₃ – Modified SRI (Application of RDF through inorganics with FYM @ 10 t ha⁻¹ and transplanting at 25 cm x 25 cm spacing)
DAT – Days after transplanting NS-Non significant

Table 2 : Grain yield and straw yield as influenced by methods of planting and age of seedlings

Treatments	Grain yield (kg ha ⁻¹)					Straw yield (kg ha ⁻¹)						
	S ₁	S ₂	S ₃	S ₄	S ₅	Mean	S ₁	S ₂	S ₃	S ₄	S ₅	Mean
M ₁	5397	5327	4997	4910	4893	5105	6433	6263	5990	5950	5903	6108
M ₂	6377	6313	6153	6143	6080	6213	7240	7216	7093	7100	7073	7144
M ₃	6440	6416	6226	6260	6370	6342	7383	7327	7143	7140	7173	7233
Mean	6071	6018	5792	5771	5781	6019	7019	6935	6742	6730	6716	7019
	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)	S.E.±	C.D. (P=0.05)
Main treatments (M)	3.9	15.3	5.1	5.1	20.2							
Sub treatments (s)	5.2	15.2	5.4	5.4	15.9							
S at same M	9.0	NS	9.4	9.4	NS							
M at same or different S	8.9	NS	9.9	9.9	NS							

M₁ – Normal method (Application of RDF through inorganics with FYM @ 10 t ha⁻¹ and transplanting at 20 cm x 10 cm spacing)
M₂ – Recommended SRI (Application of RDN through organics and transplanting at 25 cm x 25 cm spacing)
M₃ – Modified SRI (Application of RDF through inorganics with FYM @ 10 t ha⁻¹ and transplanting at 25 cm x 25 cm spacing)
DAT – Days after transplanting NS-Non significant

grain yield of rice (Table 2). Modified SRI method registered significantly higher grain yield (6342 kg ha⁻¹) which was 24.23 per cent higher over the normal method. The better performance of rice under modified SRI method may be because of favorable climatic conditions for crop growth, as could be seen through higher number of panicles per hill (61.52), panicle length (21.17 cm), number of grains per panicle (167.25) and test weight (20.83). These results are in agreement with the findings of Norman Uphoff (2002) who reported that by changing the management practices, there was increase in the grain yield under SRI method. This might be due to higher number of tillers, number of grains per panicle and panicle length. These findings are in conformity with Adrainaiivo and Jolei Barinson, (2002). Significantly lower sterility per cent was recorded in modified SRI method and which resulted in higher grain yield. Straw yield also followed same trend as that of grain yield.

Modified SRI and recommended SRI method recorded significantly higher straw yield of paddy than normal method. The increase in straw yield of paddy in modified SRI method over the recommended and normal methods was by 1.23 and 15.53 per cent, respectively. The higher straw yield under modified SRI method was due to higher dry matter production and accumulation in stem.

Age of seedlings :

The grain yield of rice was significantly influenced by age of seedlings (Table 2). Planting of 9 days (6071 kg ha⁻¹) or 12 days (6018 kg ha⁻¹) old seedlings gave significantly higher grain yield than 15 (5792 kg ha⁻¹), 18 (5771 kg ha⁻¹) and 21 days (5781 kg ha⁻¹) old seedlings. This was because of the higher number of panicle per hill (48.93 and 48. under planting of 9 days and 12 days old seedlings, respectively), panicle length (21.07 and 21.24 under planting of 9 days and 12 days old seedlings,

respectively), number of grains per panicle (159.33 and 158.13 under planting of 9 days and 12 days old seedlings, respectively) and test weight (20.74 and 20.58 g under planting of 9 days and 12 days old seedlings, respectively) and further transplanting younger seedlings *i.e.* less than 15 days old seedlings had higher tillering capacity and more vigour which inturn helped in extracting nutrients from soil. Similar findings were reported by Thiagarajan *et al.*, 2002. From this study, it could be concluded that, modified SRI method with planting of 9 or 12 days old seedlings was found to be optimum in TBP area for enhancing rice production.

REFERENCES

- Andrainaiivo, B. and Jolei Barinson (2002).** Assessment of SRI. *Proceedings of an International Conference held in Sanya, China*, April 1-4, p. 162.
- Balasubramanian, A., Venkataraman, R. and Palaniappan, S.P. (1977).** Studies on the time of planting and seedling age on the performance of rice. *Madras agric. J.*, **64**:49-51.
- Norman Uphoff (2002).** System of Rice Intensification for enhancing the productivity of land, labour and water. *J. agric. Resource Mgmt.*, **1**: 43-49.
- Pattar, P.S., Reddy, B.G.M. and Kuchanur P.H. (2001).** Yield and yield parameters of rice (*Oryza sativa* L.) as influenced by date of planting and age of seedlings. *Indian J. agric. Sci.*, **71**: 521-522.
- Thiyagarajan, T.M., Velu, V., Ramaswamy, S., Durgadevi, D., Govindarajan, K., Priyadarshini, R., Sudhalakshmi, C., Senthil Kumar, K., Nisha, P.T., Gayathry, G., Hengsdijk, H. and Bindraban, P.S. (2002).** Effects of SRI practices on hybrid rice performance in Tamil Nadu, India, In : *Water wise Rice Production*, IRRI Publication, pp. 119-127.

Received : November, 2009; Accepted : February, 2010