

Influence on the overall performance of the mulberry silkworm, *Bombyx mori* L. CSR-18 cocoon reared with V₁ mulberry leaves irrigated by distillery spentwash

S. CHANDRAJU¹, GIRIJANAGENDRASWAMY¹ AND C.S. CHIDAN KUMAR²

¹Department of Studies in Sugar Technology, Sir M. Visweswaraya Postgraduate Center, University of Mysore, Tubinakere, MANDYA (KARNATAKA) INDIA

²Department of Chemistry, Alvas Institute of Engineering Technology, Shobhavana Campus, Mijar, MOODBIDRI (KARNATAKA) INDIA

Email : chandraju1@yahoo.com

CSR-18 silkworm reared with V₁ variety of mulberry plants irrigated by raw water, 50 per cent PTSW and 33 per cent PTSW. The different parameters such as raw silk (%), filament length(m), reelability (%), denier and shell ratio were determined at the maturity of cocoons. It was found that the parameters were better in cocoon irrigated with 33 per cent PTSW compared to 50 per cent PTSW and raw water irrigations. This concludes that the mulberry plants irrigated with 33 per cent PTSW is enriched with more nutrients for the potential growth of mulberry plants which results in the potential cocoons.

Key words : Silk worm, Growth, Mulberry plant, Irrigation, Cocoon parameters

How to cite this paper : Chandraju, S., Girijagendraswamy and Kumar, C.S. Chaidanand (2012). Influence on the overall performance of the mulberry silkworm, *Bombyx mori* L. CSR-18 cocoon reared with V₁ mulberry leaves irrigated by distillery spentwash. *Asian J. Bio. Sci.*, 7 (2) : 199-202.

INTRODUCTION

The silkworm, *Bombyx mori* L. is a typical monophagous insect and mulberry (*Morus* spp.) leaf is its sole food. Man has immensely benefited from the silk produced by silkworms and subsequently researchers have always been trying to unveil the factors that can be manipulated to the benefit of the silkworm rearers. Sericulture is an age-old land-based practice in India with high employment potential and economic benefits to agrarian families. No doubt, India is the second largest producer of mulberry silk next only to China. Plants are the richest source of organic chemicals on earth and phytochemicals have been reported to influence the life and behaviour of different insects. Various extracts of medicinal plants have been tested by supplementation in the silkworm *Bombyx mori* L. were seen to influence the body weight, silk gland weight and the silk thread length in *Bombyx mori* L. (Murugan *et al.*, 1998). Dietary supplementation of the leaf, flower and pod extracts of *Moringa oleifera* and chitosan solution (Li *et al.*, 2010) elicited varied responses in the final

instar larvae of *Bombyx mori*. Nutrition plays an important role in improving the growth and development of *Bombyx mori* L. (Kanafi *et al.*, 2007). Alagumalai *et al.* (1991) observed fortification of mulberry leaves with the flour of black gram and red gram to improve the larval growth and cocoon characteristics in *Bombyx mori* L. Similarly, the growth of silkworm larvae improved significantly upon feeding them with mulberry leaves supplemented with different nutrients (Sarker, 1993). The quantity and the quality of dietary protein has long been considered to be important in the growth of the silkworm. Higher growth rate as well as weight gain can be observed in higher protein utilized group and the relative growth rate varied among the different breeds of the silkworm (Magadum *et al.*, 1996) and were influenced by the season (Isaiarasu and Suriabraman, 1999). The difference in the relative growth rate of *Aloe vera* tonic supplemented larvae from the control observed in the present study indicates that the *Aloe vera* supplementation results in higher protein utilization. Murugan *et al.* (1998) noticed a strong correlation between

the growth of silkworm and the silk production in the silkworm after the treatment with plant extracts and attributed the growth promoting effect of the plant extracts to the stimulation of biochemical processes leading to protein synthesis. The economic characters of the silk cocoon were reported to improve by feeding the silkworm with mulberry leaves treated with amino acids. It has been observed that the enrichment of mulberry leaves with calcium chloride increases the cocoon characters like cocoon weight, shell weight cocoon /shell ratio and silk proteins. The cocoon weight increased when the silkworm larvae were fed with bloodmeal fortified mulberry leaves (Matsura, 1994). Chamudeswari and Radhakrishnaiah (1994) reported the increased of cocoon weight, when the silkworm larvae were fed with zinc and nickel fortified mulberry leaves. Majumdar and Medda (1995) reported the supplementation of tyrosine to enhance the cocoon weight due to the increased synthesis of DNA, RNA and proteins in silk gland. The weight and the size of cocoon shell ratio and fibroin content of the shell increased with the supplementation of the amino acid, glycine (Isaiarasu and Ganga, 2000) reported that administration of JH analogue, methaprene, to fifth instar larvae of *Bombyx mori* L. through hypodermic injection increased the shell weight by 16 per cent over the control. Improvement in economic characters of silkworm was also noticed with folic acid administration. The silkworm larvae fed on mulberry leaves treated with *Coffea arabica* leaf extracts at 1:25 concentration recorded significantly higher.

Diluted spentwash increase the uptake of nutrients, height, growth and yield of leaves vegetables (Chandraju and Basvaraju, 2007; and Chandraju, 2008) and yields of condiments (Chidan Kumar and Chandraju, 2009), yields of some root vegetables in untreated and spentwash treated soil (Chidankumar *et al.*, 2009), yields of top vegetables (creepers) (Chidankumar *et al.*, 2009), yields of tuber/root medicinal plants (Nagendraswamy *et al.*, 2010), yields of leafy medicinal plants (Nagendraswamy *et al.*, 2010), yields of leafy medicinal plants in normal and spentwash treated soil (Chandraju *et al.*, 2010), However, no information is available on the yields of cocoon parameters of silkworms CSR-18, reared using V₁ mulberry leaves cultivated by irrigation with distillery spentwash. Therefore, the present investigation was carried out, to study the influence of V₁ mulberry leaves cultivated by irrigating with different proportions of spentwash on the cocoon parameters of silkworms CSR18, reared using V₁ mulberry leaves.

RESEARCH METHODOLOGY

Mulberry plant selected for the present study was V₁ variety. The land was ploughed repeatedly (3 to 4 times) to loosen the soil and all gravel, stones and weed were removed to get the fine soil. The ridges and furrows were made at a distance of 1.0 m, sets were planted at a distance of 0.6 m (set

to set) along the row and irrigated (by applying 5-10cm³/cm²) with raw water (RW), 50 per cent and 33 per cent SW at the dosage of once in fortnight and rest of the period with raw water (depends upon the climatic condition), without the application of any external fertilizer (either organic or inorganic). Harvesting of the leaf was done by plucking individual leaf during cooling hours of the day which was 50-60 days old. These fresh leaves were used to rear silk worms.

Disease free laying of the silkworm were obtained and raised on fresh mulberry leaves as per the new technology for silkworm rearing (Dandin *et al.*, 2000). After third moult, the larvae were acclimatized to the laboratory conditions by rearing them during the fourth instar in plastic trays of size 26 x 20 x 6 cm. During this period, they were fed four times a day. Sufficient ventilation was ensured to the larvae by placing the trays one above the other crosswise. Coolant gel bags were used to bring down the temperature and wet synthetic foam pads were used to enhance the relative humidity near the larval bed within the optimum level. A Thermo-hygrometer was used to record the temperature and relative humidity near the larval bed. Fresh and healthy leaves of V₁ variety of mulberry were used in the present study. The leaves were harvested daily from the mulberry garden during the early hours of the day and stored cool to maintain its freshness until use using wet gunny cloth in a wooden chamber. Disinfection was carried out prior to the commencement of silkworm rearing as a precautionary measure against pathogens, which may remain in the rearing room and are likely to infect the silkworm. For this, the rearing room was disinfected by spraying 2 per cent formalin solution 3 days prior to the commencement of rearing. The rearing materials such as trays and mountages were washed with chloralk solution. Dettol solution was used to wash the hands before and after handling the worms during the time of rearing. A bed disinfectant powder prepared by grinding lime powder, paraformaldehyde and benzoic acid in 97:2:1 ratio was dusted mildly on the worms daily after bed cleaning. Dead larvae if any, during the course of rearing were immediately removed and discarded properly fed with untreated mulberry leaves. Thus, the larvae in both the control and experimental trays were reared with equal quantities of leaves. The temperature and relative humidity were maintained at about 26 ± 2°C and around 70 ± 10 per cent, respectively. Several parameters were studied to assess the growth and the cocoon characteristics of *B. mori*. The mature larvae of the experimental sets were isolated and mounted on separate plastic mountage (Netrika). They were left undisturbed for four days to spin the cocoon. The cocoon were harvested. Then cocoons were collected after harvest and cleaned by removing litter. Trials were conducted thrice, cocoon parameters, such as raw silk percentage, filament length, reelability, denier and shell ratio were determined, recorded by taking the average values. These quantitative parameters were measured by the procedures

given by Sonwalkar (1993).

RESEARCH FINDINGS AND ANALYSIS

The cocoon parameters were very high reared using V₁ variety mulberry plant leaves cultivated by 33 per cent SW irrigation, and moderate in 50 per cent, while comparatively poor in RW (Table 1). In previous studies it was also found that 33 per cent SW irrigation favours the growth, yield and nutrients of plants. This could be due to the maximum absorption of NPK by the plants at 33 per cent dilution. In the case of 50 per cent SW irrigation the yields were low.

Enrichment of nutrients in V₁ mulberry leaves cultivated by 33 per cent influence healthy growth of silkworms contains

Cocoon parameters	Irrigation medium		
	RW	50% PTSW	33% PTSW
Raw silk (%)	20.00±0.010	21.50±0.011	23.40±0.013
Filament length (m)	757.65±0.009	814.34±0.012	914.66±0.009
Reelability (%)	77.80±0.012	83.33±0.017	85.20±0.008
Denier	2.62±0.017	2.69±0.010	2.78±0.012
Shell ratio	22.56±0.011	23.61±0.010	24.13±0.012

comparatively high proportion of natural protein fiber secreted by silkworms in the form a thread, fibroin–inner core comprising 75 per cent of silk, sericin -outer gum comprising 25 per cent of silk.

Conclusion :

It was observed that the parameters of cocoons produced by rearing the silk worms using V₁ variety of mulberry leaves cultivated by irrigation in 33 per cent SW were maximum and moderate in 50 per cent SW and minimum in RW irrigations. It concludes that, in 33 per cent SW irrigation the plants were able to absorb maximum amounts of nutrients (NPK) both from the soil and the spentwash resulting high yield and enhance the nutrients in plants leaves which in turn influence the better growth of silk worms containing higher proportion of silk proteins yields spinning of long silk threads in cocoons resulting in increased weight of cocoons, minimizing the cost of cultivation, and increased the parameter values of cocoons resulting in high silk production, thus elevating the economy of the farmers, since cultivation of mulberry is made without using fertilizer.

Acknowledgement :

The authors thankful to The General Manager, N. S. L. Koppa, Maddur Tq. Karnataka, for providing spentwash.

LITERATURE CITED

- Alagumalai, K., Hepshyba, C.S.S. and Ramaraj, P. (1991). Bran of pulses as extra nutrient to silkworm. *India Silk*, **30**(6) : 10-11.
- Balamurugan, R. and Isaiarasu, L. (2007). Effect of theherbal tonic “Iogen” on the growth and cocoonparameters of mulberry silkworm, *Bombyx mori* L. Proceedings of the National Seminar on Appl. Zoology ANJA College, Sivakasi (T.N.) INDIA.
- Basavaraju, H.C. and Chandraju, S. (2008). Impact of distillery spent wash on the nutrients of leaves vegetables: An investigation. *Asian J. Chem.*, **20**(7):5301-5310.
- Buenz, E.J. (2008). Aloin induces apoptosis in Jurkatcells. *Toxicol. in vitro*, **22**:422-429.
- Chamudeswari, P. and Radhakrishnaiah, K. (1994). Effectof zinc and nickel on the silkworm, *Bombyx mori* L. *Sericologia*, **34**(2) : 327-332.
- Chandraju, S. and Basavaraju, H.C. (2007). Impact of distillery spent wash on seed germination and growth of leaves vegetables: An investigation. *Sugar J. (SISSTA)*, **38**: 20-50.
- Chandraju, S., Nagendra Swamy, R., Chidan Kumar, C.S. and Nagendraswamy, Girija (2010). Influence of distillery spentwash irrigation on the yields leafy medicinal plants in normal and spentwash treated soil. *Internat. J. agric. Sci.*, **7** (1): 23-26.
- Chidankumar, C.S., Chandraju, S. and Nagendra Swamy, R. (2009). Impact of distillery spentwash irrigation on yields of top vegetables (creepers). *World Appl. Sci. J.*, **6**(9): 1270-1273.
- Chidankumar, C.S., Chandraju, S. and Nagendraswamy, R. (2009). Impact of distillery spentwash irrigation on the yields of some root vegetables in untreated and spentwash treated soil. *SISSTA*, **40**:233-236.
- Chidankumar, C.S. and Chandraju, S. (2009). Impact of distillery spentwash irrigation on the yields of some condiments: An investigation. *Sugar Tech.*, **11**(3): 303-306.
- Dandin, S.B., Jayaswal, J. and Giridhar, K. (2000). *Handbook of sericulture technologies*, Central Silk Board, Bangalore (KARNATAKA) INDIA. pp. 259.
- Isaiarasu, L. and Ganga, G. (2000). Influence of dietaryglycine supplementation on the mulberry silkworm, *Bombyx mori*. *ANJAC J.*, **17**: 47-53.

- Isaiarasu, L. and Suriabraman, S. (1999).** Seasonal differences in the biochemical composition of M₅ and MR₂ varieties grown in Sivakasi and their influence on the growth of the late age larvae of *Bombyx mori*. *J. Ecobiol.*, **11**(3) : 229-231.
- Kanafi, R.R., Ebadi, R., Mirhosseini, S.Z., Seidavi, A.R., Zolfaghari, M. and Eteban, K. (2007).** A review on nutritive effect of mulberry leaves enrichment with vitamins on economic traits and biological parameters of silkworm, *Bombyx mori* L. *Indian Sericulture J.*, **4** : 86 - 91.
- Li, Bin, Su, Ting, Chen, Xiaoling, Liu, Baoping, Zhu, Bo, Wen, Qiu, YuanFang and Xie, Guanlin (2010).** Effect of chitosan solution on the bacterial septicemia disease of *Bombyx mori* (Lepidoptera: Bombycidae) caused by *Serratia marcescens*. *Appl. Entomol. & Zool.*, **45**: 145- 152.
- Magadum, S.B., Hooli, M.A. and Magadum, V.B. (1996).** Effect of the application of juvenile hormone analogue in V instar followed by thyroxin in the pure Mysore. Breed of *Bombyx mori*. L. *Sericologia*, **32**(3):385-390.
- Majumdar A.C. and Medda, A.K. (1995).** Studies on the thyroxin and vitamin B₂ induced changes in the cycle of silkworm *Bombyx mori*. *Indian J. Physiol. & Appl. Sci.*, **29**:1-13.
- Matsura, Y. (1994).** Utilization of blood meal as the source of dietary protein for the silkworm, *Bombyx mori* L. *Japan agric. Res. Quarterly*, **28**(2):133-137.
- Murugan, K., Jeyabalan, D., Senthikumar, N., Senthilnathan, S. and Sivaprakasam, N. (1998).** Growth promoting effect of plant products on silkworm.
- Nagendraswamy, R., Chandraju, S., Nagendraswamy, Girija and Chidankumar, C.S. (2010).** Studies on the impact of irrigation of distillery spent wash on the yields of tuber/root medicinal plants. *Biomedical Pharmacol. J.*, **3**(2): 99-105.
- Nagendraswamy, R., Chandraju, S., Nagendraswamy, Girija and Chidankumar, C.S. (2010).** Studies on the impact of irrigation of distillery spent wash on the yields of leafy medicinal plants. *Nat. Env. Poll. Tech.*, **9**(4): 743-748.
- Sonwalker, T.N. (1993).** *Hand book of silk technology*, Wiley Eastern Limited, NEW DELHI (INDIA). pp.14-25.

