

# Effect of humic acid with bio-regulators on the yield of greengram [*Vigna radiata* (L.) wikzek]

K. ANANTHI<sup>1</sup>, M. KARUPPAIYA<sup>2</sup>, T. ANAND<sup>3</sup> AND M. GOMATHY<sup>4</sup>

<sup>1</sup>Department of Crop Physiology, Tamil Nadu Agricultural University, COIMBATORE (T.N.) INDIA

Email : ananthipha@yahoo.com

<sup>2</sup>Department of Agricultural Entomology, C.C.S. Haryana Agricultural University, HISAR (HARYANA) INDIA

<sup>3</sup>Cotton Research Station, VEPPANTHATTAI (T.N.) INDIA

<sup>4</sup>Maize Research Station, Vagarai, PALANI (T.N.) INDIA

Continuous adaption of cereal-based cropping system has necessitated other alternatives such as pulses, which constitute a major part in agricultural economy of India. An experiment was conducted at Millet Breeding Station, TNAU, Coimbatore during *Kharif* (July, 2006 - October, 2006) in CO 6 green gram with nine different treatments. The yield and yield components were significantly increased by the application of chemicals and bioregulators. Various yield components such as days to 50 per cent flowering, days to maturity, number of clusters per plant, number of flowers per plant, number of pods per cluster, number of pods per plant, fertility co-efficient, number of seeds per pod, pod weight per plant, seed weight per plant, seed - pod ratio, pod length, hundred seed weight, grain yield per plot, grain yield per hectare and harvest index were favorably enhanced by the foliar spray of 0.1 per cent humic acid with 0.1 ppm brassinosteroid treatment.

**Key words :** Bio regulators, Greengram, Yield

**How to cite this paper :** Ananthi, K., Karuppaiya, M., Anand, T. and Gomathy, M. (2012). Effect of humic acid with bio-regulators on the yield of greengram [*Vigna radiata* (L.) wikzek]. *Asian J. Bio. Sci.*, 7 (1) : 30 - 33.

## INTRODUCTION

Pulses occupy a strategic position in intensive as well as subsistence agriculture, as they are excellent source of dietary protein for millions of people, nutritious feed for livestock and a mini nitrogen plant having profound ameliorative effect on the soil. Green gram is extensively grown in all types of soil under varying climatic conditions. Greengram is considered as the most whole some among pulses, free from heaviness and flatulence. The green pods are used as vegetable and haulms are used as fodder. On an average, the biomass has 1.5 per cent N on fresh weight basis, serving as a good source of animal feed. Flower and pod shedding is a common problem in greengram. There is also a possibility to overcome these constraints by foliar application of bio-regulators at the pre-flowering stage, which is one of the latest trends in agriculture. These bioregulators includes humic acid (HA), naphthalene acetic acid (NAA), benzyl adenine (BA), salicylic acid (SA), brassinosteroid (BR) etc. These plant

growth regulators (PGRs) in general, help to increase the number of flowers on the plant when applied at the time of flowering. The flower and pod drop may be reduced to some extent by spraying various growth regulators on foliage (Ramesh and Thirumuguran, 2001). The foliar application of PGRs and urea significantly increased seed yield per plant (Patil *et al.*, 2005). The HA increased the yield of plants more especially, with insufficient nutrient supply than with optimum nutrient supply (Pagel, 1960). The NAA is one of the important synthetic, stable and also widely used form of auxins. Auxins had been very effective in promoting growth and development in pulse crops. Upadhyay (1994) reported that foliar spray of NAA at bud initiation and pod formation stage of chick pea increased plant height, number of branches, number of flower bud, number of flowers and yield. The main role of NAA rests with the efficient transport of sugars from the photosynthesizing parts of the plant (source) to the developing grains (sinks) and also facilitating nitrogen accumulation that probably resulted in higher total dry matter production (Kalita *et al.*, 1989). Foliar spray of 25 ppm NAA

recorded significantly higher seed yield by 21 to 22 per cent than control through increased flower production, clusters per plant, pod setting percentage and pods per plant in mungbean (Patil *et al.*, 2005). Cytokinins are adenine derivatives characterized by an ability to induce cell division in the presence of auxins. Cytokinin improves crop growth and yield through redirecting the metabolic balance of growth, delayed leaf senescence, increased photosynthetic CO<sub>2</sub> uptake and partitioning of assimilates (Brault and Maldiney, 1999). Benzyl adenine (BA) is a synthetic cytokinin, which enhanced the photosynthetic rate of the plant through the increased LAI, which ultimately resulted in higher DMA. Ramesh and Thirumurugan (2001) reported that foliar spray of 25 ppm BA registered a higher DMA in soybean. Salicylic acid belongs to the group of plant phenolics. A concentration of 100 and 500 ppm of SA affected both seed germination and seedling growth in two blackgram cultivars (*viz.*, CO 5 and T 9), but induced more flowering at 10 and 100 ppm levels, respectively (Anandhi and Ramanujam, 1997). Brassinosteroid is a novel plant growth promoting steroidal lactone which is the most biologically active compound and acting as bioregulators in crop plants. Joseph (2000) recorded an increased yield in groundnut by the foliar application of BR. Senthil *et al.* (2003) reported that foliar spray of 0.5 ppm BR increased the chlorophyll content in soybean.

## RESEARCH METHODOLOGY

A field experiment was conducted at Millet Breeding Station, TNAU, Coimbatore during *Kharif* (July, 2006 - October, 2006) in green gram (CO 6) in Randomized Block Design with nine treatments and replicated thrice. Duration of the crop was 75 days. Sowing was done on 20.07.2006 with a spacing of 30 cm x 10 cm between rows and between plants. The soil was low in available nitrogen and phosphorus content and high in potassium content. The cultural management and

plant protection measures were undertaken as and when needed. The treatment details are given below : T<sub>1</sub> - Control, T<sub>2</sub> - HA 0.1 per cent alone, T<sub>3</sub> - HA 0.1 per cent + 10 ppm benzyl adenine (BA), T<sub>4</sub> - HA 0.1 per cent + 100 ppm salicylic acid (SA), T<sub>5</sub> - HA 0.1 per cent + 0.1 ppm brassinosteroid (BR), T<sub>6</sub> - HA 0.1 per cent + micronutrient mixture, T<sub>7</sub> - HA 0.1 per cent + micronutrient mixture + 10 ppm benzyl adenine (BA), T<sub>8</sub> - HA 0.1 per cent + micronutrient mixture + 100 ppm SA and T<sub>9</sub> - HA 0.1 per cent + micronutrient mixture + 0.1 ppm BR

Sprays of 2 per cent DAP and 40 ppm NAA are common for all the above treatments. Time of sprays: First spray at peak vegetative phase and second spray at 10 days thereafter.

The important components contributing to the yield potential of the crop were recorded from the commencement of flowering to the harvest of the crop. Different yield parameters were recorded and analyzed statistically.

## RESEARCH FINDINGS AND ANALYSIS

The data collected on various pod characters as well as the yield components are presented in Table 1 and 2.

Growth regulator spray on crop growth period had significant influence on the number of clusters per plant. Foliar spray of 0.1 per cent HA + 0.1 ppm BR spray (T<sub>5</sub>) produced significantly more number of clusters per plant (7.7), which was closely followed by the foliar spray of 0.1 per cent HA + 10 ppm BA (T<sub>3</sub>) with comparable value (7.6). The treatment of NAA was effective in increasing the number of clusters per plant (Patil *et al.*, 2005) in greengram. A high correlation was also observed between number of clusters per plant and grain yield in pigeonpea (Dani, 1979). Different treatments had significant influence on the number of pods per cluster which is an important yield component. Higher number of pods per cluster was recorded with the 0.1 per cent HA + 0.1 ppm BR spray (T<sub>5</sub>) (4.1), while the minimum number was recorded with control (3.8) treatment (Table 1). Rajavel (2004) pointed out


Treatments	Days to 50% flowering	Days to maturity	No. of clusters plant <sup>-1</sup>	No. of flowers plant <sup>-1</sup>	No. of pods clusters <sup>-1</sup>	No. of pods plant <sup>-1</sup>
T <sub>1</sub>	30.6	67.5	5.7	74.6	3.8	21.6
T <sub>2</sub>	29.8	67.1	6.0	77.3	4.0	23.9
T <sub>3</sub>	27.1	65.2	7.6	94.5	4.1	31.2
T <sub>4</sub>	27.7	65.5	7.3	92.6	4.1	29.9
T <sub>5</sub>	26.3	64.7	7.7	95.5	4.1	31.7
T <sub>6</sub>	29.6	67.6	6.2	79.5	4.0	24.7
T <sub>7</sub>	28.5	66.6	6.9	89.5	4.1	28.2
T <sub>8</sub>	28.9	67.1	6.7	86.8	4.0	26.8
T <sub>9</sub>	28.1	66.3	7.4	90.5	4.1	26.6
Mean	28.5	66.4	6.8	86.7	4.0	21.65
S.E.±	0.3	0.8	0.2	1.5	0.1	1.2
C.D. (P=0.05)	0.5	1.9	0.4	3.1	0.3	2.7

**Table 2 : Effect of foliar spray of liquid humic acid with growth regulators on yield and yield components in greengram**

Treatments	No. of seeds pod <sup>-1</sup>	Pod weight plant <sup>-1</sup> (g)	Seed weight plant <sup>-1</sup> (g)	Seed / Pod ratio	Pod length (cm)	Hundred seed weight (g)	Grain yield (g plot <sup>-1</sup> )	Grain yield (kg/ha)	Harvest index (%)
T <sub>1</sub>	6.60	11.24	9.68	0.84	5.8	3.28	849	708	26.24
T <sub>2</sub>	7.28	11.61	10.58	0.86	5.7	3.52	875	729	26.88
T <sub>3</sub>	9.48	13.58	12.25	0.96	7.1	4.13	925	771	30.57
T <sub>4</sub>	8.84	13.26	11.67	0.90	6.8	4.05	912	760	30.01
T <sub>5</sub>	9.75	13.83	12.55	0.93	7.3	4.27	1047	873	31.64
T <sub>6</sub>	7.58	11.90	10.77	0.87	5.9	3.53	892	743	27.71
T <sub>7</sub>	8.57	12.51	11.30	0.89	6.5	3.85	907	756	28.85
T <sub>8</sub>	8.33	12.15	10.94	0.90	6.4	3.79	901	751	28.21
T <sub>9</sub>	8.74	12.81	11.34	0.90	6.6	3.97	912	760	29.97
Mean	8.35	12.54	11.23	0.89	6.4	3.94	916	761	29.24
S.E.±	0.14	0.47	0.48	0.03	0.10	0.28	47	57	0.99
C.D. (P=0.05)	0.30	0.99	1.02	0.06	0.19	0.59	100	120	2.13

the application of BR significantly influenced the pods per cluster resulting in higher yield in black gram. Maximum number of seeds per pod was recorded by the treatment of 0.1 per cent HA + 0.1 ppm BR (9.75), which was followed by T<sub>3</sub> (9.48) and quite comparable. Both the treatments were significantly different from control. Minimum number of seeds per pod was, however, produced by the control treatment (6.60) (Table 2). The influence of BA on number of seeds per pod was explained by Upadhyay (1994).

No significant differences were observed in seed - pod ratio due to growth regulator / nutrient sprays. However, treatment T<sub>3</sub> produced higher seed - pod ratio (0.96) which was closely followed by the spray of 0.1 per cent HA + 0.1 ppm BR (0.93). The least value was recorded by the control (0.84) (Table 2). Ghosh *et al.* (1991) pointed out that the spray of NAA at pre-flowering stage significantly increased the shelling percentage. Similar results were obtained by Gomathi (1996) in mung bean.

**Fig. 1 : Effect of foliar spray of liquid humic acid with growth regulators on grain yield (kg/ha) in greengram**

The grain yield per plot of green gram was significantly influenced by various treatments tried. Among the treatments, foliar spray of 0.1 per cent HA + BR 10 ppm (T<sub>5</sub>) ranked first with higher grain yield per plot of 1047 g plot<sup>-1</sup>, followed by T<sub>3</sub> (925). This was 23.32 and 8.95 per cent increase over the unsprayed control (849). The grain yield of greengram was significantly influenced by the foliar spray of different growth regulators and nutrients. Significantly higher yield was registered (873) with the treatment of 0.1 per cent HA + 0.1 ppm BR (T<sub>5</sub>) in comparison with the control (708). The increased yield in the best treatment was found to be 23.31 per cent over control. Second best treatment was T<sub>3</sub> (771) which was comparable with T<sub>5</sub> (Table 2 and Fig. 1). The production of higher seed yield due to growth regulators may be attributed to the fact that plants treated with growth regulators remained physiologically more active to build up sufficient food reserves for developing flowers and seeds. Thus, the plant showed early flower initiation and better seed development which ultimately resulted in higher seed yield in the treated plants. Increased grain yield was recorded by BR spray in greengram as reported by Bhatia and Jatinder Kaur (1997). Application of NAA also enhanced the number of flowers, number of pods per plant, seed yield per plant and hundred seed weight in greengram (Patil *et al.*, 2005).

## LITERATURE CITED

- Anandhi, S. and Ramanujam, M.P. (1997). Effect of salicylic acid on blackgram cultivars. *Indian J. Plant Physiol.*, **2**: 138-141.
- Bhatia, D.S. and Kaur, Jatinder (1997). Effect of homobrassinolide and humicil on chlorophyll content, hill activity and yield components in mungbean [*Vigna radiata* (L.) Wilczek]. *Phytomorphol.*, **47**: 421-426.

- Brault, M. and Maldiney, R. (1999).** Mechanism of cytokinin action. *Plant Physiol. Biochem.*, **37**: 403-412.
- Dani, R.G. (1979).** Variability and association between yield and yield components in pigeonpea. *Indian J. Agric.Sci.*, **49**: 507-510.
- Ghosh, R.K., Mandai, Bikash Kumar and Chatterjee, B.N. (1991).** Effect of growth regulators on the productivity of some major oilseed crops. *J. Agron. Crop Sci.*, **167**: 221-228.
- Gomathi, R. (1996).** Chemical manipulation on yield improvement in greengram [*Vigna radiata* (L.) wilczek] M.Sc. (Ag.) Thesis, Tamil Nadu Agricultural University, COIMBATORE, T.N. (India).
- Joseph, Bindhu (2000).** Physiological and biochemical effects of brassinolide on the productivity of groundnut. M.Sc. (Ag.) Thesis, Tamil Nadu Agricultural University, COIMBATORE, T.N. (India).
- Kalita, M.M. (1989).** Effect of phosphate and growth regulators on greengram. *Indian J. Agron.*, **34**: 236-237.
- Pagel, H. (1960).** Über den Einfluss Von Humusstoffen auf das Pflanzenwachstum. I. Einfluss. Von Humusstoffen auf Keimung und Wurzelwachstum. *Albrecht. Thier. Archiv.*, **4**: 450-468.
- Patil, S.N., Patil, R.B. and Suyawanshi, Y.B. (2005).** Effect of foliar application of plant growth regulators and nutrients on seed yield and quality attributes of mungbean [*Vigna radiata* (L.) Wilczek]. *Seed Res.*, **33**: 142-145.
- Ramesh, K. and Thirumurugan, V. (2001).** Effect of seed pelleting and foliar nutrition on growth of soybean. *Madras Agric. J.*, **88**: 465-468.
- Rajavel, S. (2004).** Physiological studies on increasing yield potential in blackgram [*Vigna mungo* (L.) Hepper] genotypes with plant growth regulators. M.Sc. (Ag.) Thesis, Tamil Nadu Agricultural University, COIMBATORE, T.N. (India).
- Senthil, A., Pathmanaban, G. and Srinivasan, P.S. (2003).** Effect of Bioregulator of some physiological and biochemical parameters of soybean (*G. max* L.). *Ley Res.*, **28**: 54-58.
- Upadhyay, R.G. (1994).** Effect of bioregulators on growth, development, flowering behaviour and yield of chickpea (*Cicer arietinum* L.). *Legume Res.*, **17**: 60-62.

---

Received : 01.12.2011; Revised : 05.01.2012; Accepted : 01.02.2012