

Relationship of waist-hip ratio and body mass index to blood pressure among adult female students

■ JIGMAT DACHEN AND UJWALA KOCHÉ¹

Received : 21.03.2014; Revised : 07.09.2014; Accepted : 20.09.2014

Members of the Research Forum

Associated Authors:

¹Dr. Ambedkar College,
CHANDARAPUR (M.S.) INDIA

Author for correspondence :

JIGMAT DACHEN

Directorate of Physical Education and
Sports, University of Kashmir,
KASHMIR (J&K) INDIA
Email: imjigi@yahoo.co.in

■ ABSTRACT

The relationship between waist - hip ratio (WHR) and body mass index (BMI) with blood pressure has not been much studied among healthy female in India. Therefore, this study was undertaken to test the relative efficacy of waist hip ratio (WHR) and body mass index (BMI) to predict blood pressure among healthy college going female. Thirty one (31) healthy female college students who participate in regular physical activities were tested on height, weight, hip circumference, waist circumference, systolic blood pressure and diastolic blood pressure. Standard anthropometric techniques and formula were used. Pearson product moment correlation and multiple regression analysis were employed to analysis the data. There was significant relationship ($p=.05$) between body mass index (BMI) with systolic blood pressure ($r=.42$) and with diastolic blood pressure ($r=.45$). Further regression analysis revealed that body mass index (BMI) explained 15 per cent of variance of systolic blood pressure and 18 per cent of variance of diastolic blood pressure among healthy college going female. However waist hip ratio (WHR) has not found significant relation with systolic and diastolic blood pressure.

■ Key Words : Waist-hip ratio, Body mass index, Blood pressure

■ How to cite this paper : Dachen, Jigmat and Koche, Ujwala (2014). Relationship of waist-hip ratio and body mass index to blood pressure among adult female students. *Internat. J. Phy. Edu.*, 7 (2) : 59-62.

The World Health Organisation reports that the average blood pressure went down by 2.7mm mercury among women globally, while increasing by 2.4mm mercury in India. In men, it decreased by 2.3 mm mercury globally in the past three decades but in India it went up by 2.2 mm. mercury. High blood pressure is the leading risk factor for cardiovascular disease mortality, causing more than 7 million deaths every year worldwide. The Harvard study presented at the World Congress of Cardiology in Dubai says around 900 million people in developing countries have high blood pressure but only one-third are aware of their disease. Moreover, only 100 million of these people receive treatment, while only 5 per cent of the total is controlled. High blood pressure is twice as common in adult who are obese than those who are healthy (Times of India, Date: 22/04/2012).

Obesity is associated with high triglycerides and decrease HDL cholesterol.

Person with central abdominal obesity, characterised by excessive visceral fat and abdomen have higher risk of weight related disease. Increased body fat is accompanied by profound changes in the physiological and metabolic functions of the body, which are directly dependent on the degree of excess weight and on its distribution around the body. Body mass index has been identified by the World Health Organization (WHO) as the most useful epidemiological measure of obesity. However, it does not take into account the distribution of body fat, resulting in variability in different individuals and populations (WHO, 2000). Hence, other anthropometric indices like waist circumference and hip measures the central distribution of fat. Waist hip ratio has been recommended as

a simple and practical measure for identifying overweight and obese patients. It is particularly useful for individuals and population groups with different body builds (Larson *et al.*, 1984; Lapidus *et al.*, 1984; Welborn *et al.*, 2003). Seidell *et al.* (1989) have suggested that BMI was the best overall predictor for both systolic and diastolic blood pressure for women. However Enrique Rivero (2009) study revealed that waist-hip ratio better than BMI for gauging obesity in elderly. Therefore present study is attempted to identify relationship of body mass index (BMI) and waist-hip ratio (WHR) with systolic and diastolic blood pressure among healthy college going female subjects.

METHODOLOGY

The present investigation was undertaken to determine the waist-hip ratio and body mass index to blood pressure among adult students. The following procedure and statistical analysis has been used :

Participants :

Thirty one (31) healthy female college students (Mean age 21.13±2.05) who participate in regular physical activities were tested on height, weight, hip circumference, waist circumference, systolic blood pressure and diastolic blood pressure. Standard anthropometric techniques and formula were used to determine the body mass index (BMI) and waist-hip ratio (WHR). The Selection was by a random sampling with the co-operation and readiness of the participants after their informed consent.

Instrumentations :

Standard instruments like Gulick Tape, Stadiometre, Sphygmomanometer, and portable weighing scale were used by female researcher to collect the data.

Statistical analyses :

All data were analyzed by SPSS (Statistical Package for social sciences, Version 17, SPSS). Mean, standard deviation, Pearson’s correlation test, regression analysis and ANOVA were used to investigate the relationship between the body mass index (BMI) and waist-hip ratio (WHR) and systolic and diastolic blood pressure.

OBSERVATIONS AND DISCUSSION

Table 1 presents the Mean and Standard of body mass index (BMI), waist-hip ratio (WHR), systolic blood pressure and diastolic blood pressure. All the parameters are of normal range as per world health organisation (WHO) norms.

Table 2 shows body mass index (BMI) has significant relationship with systolic blood pressure (r=.42) and with diastolic blood pressure (r=.45). However there is no significant relationship between waist-hip ratio with systolic blood pressure and diastolic blood pressure. Significant relationship was found between body mass index (BMI) and waist-hip ratio (WHR), and between systolic blood pressure and diastolic blood pressure (r=.54), and (r=.59), respectively.

Table 3 present model summary of regression analysis which showed body mass index explained the 15% of variance (as adjusted R Square value is .05) of systolic blood pressure,

Table 1 : Descriptive statistics

Variables	Mean	Std. deviation
Body mass index	22.18	2.90
Waist hip ratio	0.80	0.04
Systolic blood pressure	107.83	6.33
Diastolic blood pressure	70.38	6.67

N=31

Table 2 : Correlation matrix among Independent and dependent variables

	Body mass index	Waist hip ratio	Systolic blood pressure	Diastolic blood pressure
Body mass index	1			
Waist hip ratio	0.54*	1		
Systolic blood pressure	0.42*	0.09	1	
Diastolic blood pressure	0.45*	0.00	0.59*	1

N=31, Significant at 0.05 Level

Table 3 : Regression model summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	0.42a	0.17	0.15	5.83

Predictors : (Constant), body mass index

Dependent variable : systolic blood pressure




Fig. 1 : Showed the scatter graph of body mass index (BMI) with systolic blood pressure, linear relationship was found between this two variables




Fig. 2 : Showed the scatter graph of body mass index (BMI) with diastolic blood pressure, linear relationship was found between this two variables

and this model is significant at .05 level, as F value= 6.3 (ANOVA) was significant at .05 level.

Table 4 present model summary of regression analysis which showed body mass index explained the 17 per cent of variance (as adjusted R Square value is .17) of systolic blood pressure, and this model is significant at .05 level, as F value 6.3 was significant.

Finally two regression equation has been develop in order to predict the cardio vascular risk factor *i.e.*, systolic blood pressure and diastolic blood pressure. Where Y_1 and Y_2 are the predicted systolic and diastolic blood pressures and 87.42 and 47.19 are predictors constant, X_1 and X_2 are raw scores of Body mass index (B.M. I) of subjects and 0.93 and 1.05 are unstandardized B coefficients value for equation 1 and 2, respectively.

$$Y_1 = 87.42 + (X_1) \times 0.93$$

$$Y_2 = 47.19 + (X_2) \times 1.05$$

The more obese a person is, the more likely he or she is to develop health problems. Mild obesity involving a body mass index (BMI) of 30+ is less dangerous to health than morbid obesity (BMI 40+) or malignant obesity (BMI 50+). Obesity has been particularly recognized as a major independent risk factor for cardiovascular diseases (Despres *et al.*, 2001). This is because increased body fat is accompanied by profound changes in the physiological and metabolic

functions of the body, which are directly dependent on the degree of excess weight and on its distribution around the body.

The prevalence of obesity is rising in developed and developing nations, and it is cited as an important risk factor for early mortality (WHO, 1998). Obesity has a strong relationship with cardiovascular diseases like hypertension (Stevens *et al.*, 1998; Despres *et al.*, 2001 and Cameron *et al.*, 2003), coronary heart disease and diabetes. A number of clinical measurements for obesity have been used to determine susceptibility to cardiovascular diseases (Cameron *et al.*, 2003). These include anthropometric indices such as body mass index (BMI), waist-hip ratio (WHR) and waist circumference (WC) (Bray and Gray, 1988; Flier and Maratos, 2005).

In this study, the mean values of body mass index (BMI) and waist-hip ratio (WHR) for female (BMI value=22.18 and WHR value =0.80) participants fall within the normal range and classified as average fat under world health organisation (WHO) norms. In addition, the result of the study showed that body mass index (BMI) has significant correlation with systolic blood pressure and diastolic blood pressure and strong independent indicator of blood pressure. However waist-hip ratio is not found significant relationship with systolic and diastolic blood pressure, and this study is partially in accordance with Canoy *et al.* (2004); Dowling and Pi-Sunyer (1993); Nesto (2003); (Flier and Maratos, 2005) where they

Table 4 : Regression model summary				
Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.45a	.20	.18	6.04

Predictors: (Constant), body mass index
 Dependent variable: diastolic blood pressure

found BMI and WHR, correlated significantly with systolic and diastolic blood pressures. Finally regression equation is being developed in order to predict the systolic blood pressure and diastolic blood pressure.

Acknowledgment :

Authors are very grateful to Dr. Anil Kumar Karwande, Principal, Ishwar Deshmukh College of Physical Education, Nagpur for giving permission to conduct the study.

■ REFERENCES

- Bray, G.A. and Gray, D.S. (1988).** Obesity. Part 1- Pathogenesis. *West J. Medicine*, **149** (4) : 429-441.
- Cameron, A.J., Welborn, T.A. and Zimmet, P.Z. (2003).** Overweight and obesity in Australia: The 1999-2000 Australian Diabetes, Obesity and Lifestyle Study. *Medical J. Australia*, **178** (9) : 427-432.
- Canoy, D., Luben, R., Welch, A. (2004).** Abdominal obesity and respiratory function in men and women in the EPIC-Norfolk Study, United Kingdom. *Am. J. Epidemiol.*, **159**: 1140-1149.
- Despres, J.P., Lemieux, I. and Prud'Homme, D. (2001).** Treatment of obesity, need to focus on high risk abdominally obese patients. *British Medical J.*, **322** (7288) : 716-720.
- Dowling, H.J. and Pi-Sunyer, F.X. (1993).** Race- dependent health risks of upper body obesity. *Diabetes*, **42** (4) : 537-543.
- Flier, J.S. and Maratos, E. (2005).** Obesity. In: Kasper, D.L. and Fauci, A.S. (eds). *Harrison's Principles of Internal Medicine* 16th ed. New York, McGraw-Hill, pp.423-40.
- Larsson, B., Svardsudd, K. and Welin, L. (1984).** Abdominal adipose tissue distribution, obesity and risk of cardiovascular disease and death: 13 year follow-up of participants in the study of men born in 1913. *Br. Med. J. (Clin. Res. Ed.)*, **288**(6428): 1401-1404.
- Nesto, R.W. (2003).** The relation of insulin resistance syndromesto risk of cardiovascular disease. *Rev. Cardiovascular Medicine*, **4**(6): 11-18.
- Seidell, J.C., Cigolini, M., Charzavrka, J., Elsinger, B.M.D., Biase, G., Bjorntosp, P., Hautvast, J.G.A.J., Contardo, F.S., Zostak, V., and Scuro, L.A. (1989).** Indicators of fat distributions, Serun lipids, and blood pressure in European women born in 1948- the European fat distribution study. *Am. J. Epidemiol.*, **130**: 53-55.
- Stevens, Bjorn, William R. Cotton, Graham Feingold and Moeng, C.H. (1998).** Large-eddy simulations of strongly precipitating, shallow stratocumulus-topped boundary layers. *J. Atmos. Sci.*, **55**: 3616-3638.
- Welborn, T.A., Satvinders, D. and Bennet, S.A. (2003).** Waistship-ratio is the dominant risk factor predicting cardiovascular death in Australia. *Medical J. Australia*, **179** (11-12) : 580-585.
- World Health Organization (1998).** Obesity: preventing and managing the global epidemic. Report of a WHO consultation on obesity.

■ WEBLIOGRAPHY

World Health Organization (2000). Obesity: preventing and managing the global epidemic. Geneva (WHO Technical Report Series, No. 894). http://whqlibdoc.who.int/trs/WHO_TRS_894.pdf.

7th
Year
★★★★ of Excellence ★★★★★