

Community structure of clambeds in tropical estuary, Mangalore

■ N. SIVASHANKAR AND GANGADHARA GOWDA

Article Chronicle :

Received :

01.11.2013;

Revised :

15.11.2013;

Accepted :

30.11.2013

SUMMARY : A study was conducted to know the community structure of the clam beds of Mulky estuary in Mangalore. The seasonal and spatial variations in the qualitative and quantitative composition of the macrobenthic population indicated the dominance of mollusca, polychaetes and crustaceans. Higher populations were observed during the premonsoon and post monsoon season. The population density of macrobenthos varied between 228 to 915 no/m². Four species of polychaetes were recorded in clam beds in which *Seballaria* sp. and *Dendroneries* were the most abundant. Crustaceans were represented by amphipods, isopods, barnacles and crabs. Molluscan population were represented by bivalves, and gastropods. Four species of bivalves were recorded in Mulki estuary among which, *Meretrix casta* and *Paphia malabarica* were the dominant. Bivalves contributed significantly to the macrobenthic population.

HOW TO CITE THIS ARTICLE : Sivashankar, N. and Gowda, Gangadhara (2013). Community structure of clambeds in tropical estuary, Mangalore. *Asian J. Environ. Sci.*, 8(2): 125-129.

Key Words :

Clam beds, Macro benthos, Monsoon seasons

Bivalve molluscs such as oysters, sea scallops, clams, and soft-shell clams, often aggregate in dense groupings, forming a habitat type known as shellfish beds. Shellfish beds provide many habitat services including refuge for smaller organisms such as polychaete worms, juvenile crabs, snails, and sea stars. Additionally, organisms such as slipper shells, sponges, and algae can attach to shells' hard surfaces in areas otherwise dominated by soft sediments. Shellfish beds provide many important ecosystem services.

The concepts of benthic community were expanded by Petersen (1913, 1918), Mollander (1930) and Thorson (1955a, 1957). They suggested that a community is any assemblages of species and individuals, as an organized unit that species composition and their frequency show regular changes in time and space, and that these numerical components keep a certain balance through a definite interspecific food relationship.

Because of environmental complexity in the habitat, benthos in the shallow waters show various living manners, though all of them occupy a position at or near the end of a food chain items

in marine ecosystems. Benthos play an important role in the recycle or regeneration of nutrients between pelagic and benthic divisions, that is, without such larger animals as megalobenthos they form an important source of the demersal fishes which work an indispensable role, in trophic levels and also provide food directly or indirectly through the detritus food chains for various micro consumers in a lower trophic level. Thus, benthos with a role of this kind have a certain domain as a habitat according to their proper manners and form a community in which certain species will occur together in a prescribed areas and the response of the benthic communities to the environmental conditions in the gradient will give not only the important clue to study the characteristics of the organic matter flow in the bottom area (Sbigeru 1982). Hence this study has been carried out to know the community structure of the clam bed in Mulky estuary.

EXPERIMENTAL METHODOLOGY

The confluence of the Mulki and Pavanje rivers with the Arabian Sea leads to the formation

Author for correspondence :

N. SIVASHANKAR

College of Agriculture,
BHEEMAREAYANAGUDI
(KARNATAKA) INDIA
Email: sivareddyn@rediffmail.
com

See end of the article for
Copied authors'

of Mulki estuary. The Mulki estuary (Lat. 13° 4' N and Long. 74° 17' E) is situated at about 45 km north of Mangalore. The estuary is connected to the sea throughout the year and is subjected to tidal influence to the tune of 1.86 m affecting the water to a length of 6.0 km in Mulki river. The bottom is generally sandy with muddy stretches in the deeper areas. Four stations have been selected in the predetermined clambeds (Fig. A).



In Mulki estuary, the clam shell deposits extend up to a depth varying between 0.5 and 1 foot below the live clam bed level. The shell deposits are of recent formation and are not fossilized. The clam beds occur in areas at a water depth of 0.5- 1.5 m during low tide where live clams formed 57% of the shell layer covering an area of 0.27 sq.km.

Samples were collected at fortnightly intervals (August, 2007-July, 2008) from the selected stations in Mulki estuary. The samples were collected during low tide by using a quadrant having an area of 0.16m², upto a depth of 10 cm and were used for further analysis. For collection of macrobenthos - the associated macrofauna, sediment samples were sieved through a 0.5mm. mesh size sieve and the fauna retained in the sieve was preserved in 5% buffered formalin with Rose Bengal solution for further analysis on the distribution, population (no./m²) and species composition of benthic fauna associated with the clambeds.

EXPERIMENTAL FINDINGS AND DISCUSSION

In the present investigation, three major groups were recorded which belonged to mainly to molluscs, polychaetes and crustaceans and were found more frequently with greater numbers. The other organisms occurred sporadically.

The numerical abundance of macrobenthic organisms varied between stations and months. The percentage

contribution of polychaetes varied from 3.02 to 21.28 per cent, whereas crustaceans varied from 1.05 to 3.22 per cent. The molluscs were most dominant and represented by gastropods, bivalves. The gastropods percentage contribution varied from 3.17 to 41.17 per cent in the clambeds the percentage contribution of bivalve varied from 38.87 to 58.68 per cent. The other miscellaneous forms varied from 4.44 to 5.97 per cent.

The percentage contribution of macrobenthos in the clambeds were found to be varied in space and time Fig 1. During monsoon season, bivalves and gastropods formed 44 and 26% of the total macro benthos, respectively. Between different months bivalves were recorded not less than 30 per cent in almost all the months followed by gastropods. In post-and premonsoon seasons similar trend was noticed. Bivalve species, *Meritrix casta* and *Paphia malabarica* were the dominant one. In gastropods, the dominant group was *Cerithidea fluviatilis* in almost all the months. The percentage contribution of spats (*M. casta* and *P. malabarica*) which were the third contributor of benthic population varied between 3 and 12 per cent. The benthos in the clambeds were dominated by bivalves, gastropods and spats (34.14 to 37.16; 24.86 to 27.80 and 14.42 to 22.54 %), respectively (Fig. 1).

The contribution of polychaetes among the seasons varied from 1 to 5 % in monsoon season; 2 to 18% in post monsoon season and 0 to 1 % in pre monsoon season, respectively. The most dominating species of polychaetes in the clambeds were *Dendronereis arborifera*.

In the backwaters and estuaries of Cochin and Vembanad lake along the west coast of India, observed the dominance of polychaetes in the benthic community (Ansari 1974, Kurian *et al.* 1975, Kurian 1971, Pillai 1977, Ansari 1977, Sunil Kumar 1995, Kumar 1997. Sunilkumar 2002). Ramachandra *et al.* (1984) recorded higher number of *Dendronereis arborifera* and *Sabellaria cementarium* in the polychaete community of Mulki estuary along west coast of India. In the present study also, similar results have been noticed.

The dominant species crustaceans observed were *Corophium sp* and *Gammarids*. The barnacles observed during the period were found to be settled on small pebbles and shells of molluscs. The monthly distribution among the stations was zero to 21.88 % with a mean of 34.80±9.77 no/m².

Studies carried in the estuaries of Mandovi-Zuari, Coleron indicated the dominance of crustaceans during post monsoon.(Ansari *et al.* 1986, Prabhadevi 1994). However er, Prabhu *et al.* (1993), Mohan Kumar (1999) and Shanthanagouda (2001) did not observe significant contribution of crustaceans in general and amphipods in particular to the total macrobenthos in the Gangolli, Mangalore coastal waters and Nethravati-Gurupur estuaries, respectively. In the present study also, the contribution of




Fig. 1 : Percentage distribution of different macrobenthos in the clam beds

crustaceans was negligible to the total macrobenthos in Mulki estuary.

The gastropod percentage in the clambeds ranged from

24 to 32% during the monsoon season; 22 to 28% in premonsoon months and 22 to 28% in post monsoon months. The most dominant species are *Cerithidea*

Table 1: Correlation co-efficients of total fauna in relation to different physico-chemical parameters in the clambeds

Populations	Water temp	Sediment temperature	Water pH	Sediment pH	Salinity	Interstitial salinity	Dissolved oxygen	Ammonia	Chlorophyll	Susp. solids	Organic carbon	plankton	Sand	Silt	Clay	V. Coarse sand	Coarse sand	Medium sand	Fine sand	V. Fine sand
Total benthos	0.425	0.551	0.671	0.630	0.717	0.749	0.678	-0.614	0.447	0.749	-	-	-	-	-	-	-	-	0.481	1.000
Polychaetes	-	-	-	-	0.398	0.445	-	-	-	0.445	-	-	-0.506	0.000	0.430	-	-	-	0.562	0.545
Crustaceans	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Bivalves	-	-	0.412	-	-	-	0.410	-	0.384	-	-	-	-	-	-	-	-	-	-	0.405
Total spats	-	-	-	0.421	0.398	0.419	0.328	-	-	0.419	-	-	-	0.455	-	-	-	-	-	0.612
Gastropods	-	0.328	0.405	0.435	0.457	0.447	0.382	-0.381	0.363	0.447	-	-	-	-	-	-	-	-	-	0.609
Miscellaneous	-	0.397	-	-	-	0.391	-	-0.460	0.405	0.391	-	-	-	-	-	-	-	-	-	-
M.c. spats	0.381	0.423	0.436	0.562	0.511	0.542	0.431	-0.428	0.323	0.542	-	-	-	-	-	-	-	-	-	0.741
P.m. spats	0.347	0.420	0.448	0.542	0.516	0.537	0.460	-0.405	-	0.537	-	-	-	-	-	-	-	-	-	0.796
M. casta	-	-	-	-	-	-	-	-	0.305	-	-	-	0.637	-	-0.614	-	0.545	-	-	0.183
P. malabarica	-	-	-	-	-	-	-	-	-	-	-	-	-0.729	-	0.671	-0.504	0.053	0.464	0.082	

fluviatilis, Neritina oualaniensis Nerita pollita and Eunaticina papilla Studies conducted in cochin backwaters, Mulki estuary, Kali estuary indicated the higher contribution of mollusks to the total macrobentic community in the calm beds (Ansari, 1977; Ramachandra et al., 1984 and Bhat and Neelakantan, 1988)

The bivalve population consisted of spats and the adult clams. The clams were mainly dominated by only two species viz., Meretrix casta and Paphia malabarica in all the stations. The percentage contribution of clams towards total population of varied from 47.93 to 67.71% during monsoon season, 34 to 48 % in post monsoon season and 34 to 37 % in pre monsoon season.

Ramachandra et al. (1984) documented low number of molluscs during monsoon and high population during post and pre monsoon in Mulki estuary. Devassy et al. (1987), Mohan Kumar (1999) observed the peak abundance of molluscs during post monsoon season. Shanthanagouda (2001) observed dominance of molluscs during post and pre monsoon season. In the present study also, similar trend was noticed, viz dominance mollusca in the post and pre monsoon season.

Sand tubes, mud tubes and egg cases formed the bulk of miscellaneous forms. During monsoon season the density of these varied from 3 to 4 no/m² in monsoon season; 5 to 10 in post monsoon season and 3 to 10 no/m² pre monsoon season, respectively.

Correlation studies indicated that the polychetes had positive relationship with san, clay and very fine sand indicating its preference. A positive correlation was observed with respect to gastropods with all the parameters except ammonia. The molluscs showed positive correlation with parameters like pH, dissolved oxygen chlorophyll and with very fine sand (Table 1).

Conclusion :

It can be concluded that, the community of clam bed were comprised of Polychetes, Gastropods, Molluscs, Crustaceans, Spats of clams viz., Meretrix casta, Paphia malabarica.

Coopted Authors' :

GANGADHARA GOWDA, Department of Fisheries Environment and Ecology, College of Fisheries, MANGALORE (KARNATAKA) INDIA

REFERENCES

Ansari, Z.A. (1974). Macrobenthic production in Vembanad lake. *Mahasagar*, 6(3-4) : 197-200.
 Ansari, Z.A. (1977). Macrobenthos of the Cochin backwaters. *Mahasagar*, 10(3-4) : 169-171.

- Ansari, Z.A.**, Ingole, B.S., Banerjee, G. and Parulekar, A.M. (1986). Spatial and temporal changes in benthic macrofauna from Mandovi and Zuari estuaries of Goa, west coast of India. *Indian J. Mar. Sci.*, **15**: 223-229.
- Bhat, U.G.** and Neelakantan, B. (1988). Environmental impact on the macrobenthos distribution of Kali estuary, Karwar, Central west coast of India. *Indian J. Mar. Sci.*, **17**(2) : 134-142.
- Kurian, C.V.** (1972). Ecology of benthos in the tropical estuary. *Proc. Indian Nat. Acad.* **38B** (3)&(4): 154-163.
- Kurian, C.V.**, Damodaran, R. and Antony, R. (1975). Bottom fauna of the Vembanad lake. In: Kurian C.V. (Ed.), Contribution to estuarine biology, *Dept. Mar. Sci. Univ.; Cochin*. pp. 987-994.
- Kumar, R.S.** (1997). Vertical distribution and abundance of macro-invertebrates in an estuarine mangrove biotope, southwest coast of India. *Indian J. Mar. Sci.*, **26**(1): 20-25.
- Mohan Kumar, B.** (1999). Distribution of macrobenthic organisms in relation to the selected sediment characteristics in the Arabian Sea off Chitrapur receiving industrial effluents. M.F.Sc., Thesis, University of Agricultural Sciences, BENGALURU, KARNATAKA (INDIA).
- Molander, A.R.** (1930). Animal communities on soft bottom areas in the Gullmar Fjord. *Kristinebergs Zoologiska Sta.*, **1877-1927** (2): 1-90.
- Nakao, Sbigeru** (1982). Community structures of the macrobenthos in the shallow waters in northern japan. *Memoirs of the Faculty of Fisheries Fokkaido University*, **28**(2) : 225-304.
- Petersen, C.G. Joh.** (1913). Valuation of the sea, II. The animal communities of the sea bottom and Their importance for marine zoogeography. *Rep. Danish Biol. Stat.*, **21**: 1-44.
- Petersen, C.G. Joh.** (1918). The sea bottom and its production of fish-food. A survey of the work done in connection with the valuation of the Danish waters from, 1883-1917. *Ibid.*, **25** : 1-62.
- Prabhadevi, L.** (1994). Ecology of Coleroon estuary: studies on benthic fauna. *J. Mar. Biol. Assoc. India.*, **36**(1)&(2): 260-266.
- Prabhu, H.V.**, Narayana, A.C. and Katti, R.J. (1993). Macrobenthic fauna in nearshore sediments off Gangolli, west coast of India. *Indian J. Mar. Sci.*, **22**: 168-171.
- Ramachandra, U.**, Gupta, T.R.C. and Katti, R.J. (1984). Macrobenthos and sediment characteristics of Mulki estuary, west coast of India. *Indian J. Mar. Sci.*, **13** : 109-112.
- Shanthanagouda, A.H.** (2001). Macrobenthos in relation to sediment characteristics in Nethravati-Gurupur estuary. M.F.Sc. Thesis, University of Agricultural Sciences, BENGALURU, KARNATAKA (INDIA).
- Sunil Kumar, R.** (1995). New record of five Annelids (Class: Polychaeta) from the mangrove habitat of the south west coast of India. *J. Mar. Biol. Assoc. India.*, **41**(1&2): 116-118.
- Sunil Kumar, R.** (2002). Biomass, horizontal zonation and vertical stratification of polychaete fauna in the littoral sediment of Cochin estuarine mangrove habitat, south west coast of India. *Indian J. Mar. Sci.*, **31**(2): 100-107.
- Thorson, G.** (1955a). Modern aspects of marine level-bottom animal communities. *J. Mar. Res.*, **14** : 387-397.
- Thorson, G.** (1957). Bottom communities (sublittoral or shallow shelf). *Gwl. Soc. Amer. Mem.*, **67** (Treat. Mar. Ecol. Palaeoecol., vol. 1. Ecology, ed. by Hedgpeth, J.W.), 461-534.


 ★★★★★of Excellence★★★★★