

Effect of combined application of composted pressmud, coirpith and farmyard manure on the yield and growth characteristics of green gram (*Vigna radiata* L.)

■ ANJU SINGH AND A.VIJAYALAKSHMI

SUMMARY

A field experiment was carried out at Avinashilingam Institute for Home Science and Higher Education for Women, Coimbatore for assessing the effect of composted coirpith, composted pressmud, farmyard manure and NPK on vegetative and yield parameters of green gram (*Vignaradiata* L.). On 25th day after sowing a significant increase in root length in T₂(composted coirpith), shoot length in T₅(NPK 100%), number of leaves in T₇(composted pressmud+50%NPK), number of nodules in T₄(FYM), fresh weight in T₁₀(composted pressmud+25%NPK) and dry weight in T₁₁(FYM + 25%NPK) was noted. On 45th day an increase in root length in T₁₀(composted pressmud+25%NPK), shoot length in T₆(composted coirpith + 50%NPK), number of leaves in T₃(composted pressmud), number of nodules in T₄(FYM), number of flowers in T₈(FYM + 50%NPK), fresh weight in T₈(FYM + 50%NPK), dry weight in T₈(FYM + 50%NPK) was observed. On 55th day a significant increase in root length in T₅(NPK 100%), shoot length in T₃(composted pressmud), number of nodules in T₃(composted pressmud), number of fruits in T₆(composted coirpith + 50%NPK), fresh weight and dry weight in T₁₂(composted coirpith + composted pressmud + FYM) was observed. There was an increase in yield parameters number of pods/plant was increase in T₁₁(FYM +25%NPK), length of pods in T₁₀(composted pressmud + 25%NPK), weight of pods in T₁₀(composted pressmud+25%NPK), number of seeds / pod in T₅(NPK 100%), weight of seeds/pod in T₈(FYM + 50%NPK), pods fresh weight in T₁₀(composted pressmud +25%NPK) and pods dry weight in T₂(composted coirpith). The positive effect obtained from composted coirpith, composted pressmud and farmyard manure in this study favour the recycling of agrowastes for sustainable crop production.

Key Words : Green gram, Composted coirpith, Composted pressmud, Farm yard manure

How to cite this article : Singh, Anju and Vijayalakshmi, A. (2013). Effect of combined application of composted pressmud, coirpith and farmyard manure on the yield and growth characteristics of green gram (*Vigna radiata* L.). *Internat. J. Plant Sci.*, 8 (2) : 410-413.

Article chronicle : Received : 11.12.2012; **Revised :** 03.06.2013; **Accepted :** 21.06.2013

Green gram (*Vigna radiata* L.) commonly known as “mung bean” is one of the most important short duration pulse in India. Among the pulse green gram occupies 10.2 lakh hectares (4.3% of total) in India. Recycling of agro industrial wastes (press mud and coir pith) can bring

MEMBERS OF THE RESEARCH FORUM

Author to be contacted :

A. VIJAYALAKSHMI, Department of Botany, Avinashilingam Institute for Home Science and Higher Education for Women, COIMBATORE (T.N.) INDIA
Email: a.vijayalakshmi85@gmail.com

Address of the Co-authors:

ANJU SINGH, Department of Botany, Avinashilingam Institute for Home Science and Higher Education for Women, COIMBATORE (T.N.) INDIA

tremendous benefits to agriculture and land management. It also gives a cleaner environment, a healthier habitat and an intelligent use of wastes. Treatment and reutilization of the agro industrial wastes in an effective way is an urgent need. The need of the hour is to improve soil health by providing the much needed organic matter. The use of organic matter and compost enhances soil organic carbon more than application of the same amount of nutrients as inorganic fertilizers.

MATERIAL AND METHODS

The agro-industrial waste such as pressmud was collected from Bannari Sugars Private Limited Sathyamangalam and coirpith from Pollachi. Seeds of green gram and FYM were collected

from Tamil Nadu Agricultural University, Coimbatore. Using *Pleurotus Sajor Caju* the compost was prepared. The pots were filled with 7kg of sandy clay loam soil. The composts were applied to the respective pots and mixed thoroughly. Viable seeds were selected and five seeds were sown in each pot with three replications. After germination three healthy plants were maintained per pots. In this experiment composted coirpith, composted pressmud and FYM were incorporated in different concentration -T₁ - Control, T₂ - Composted coirpith (12.5t ha⁻¹), T₃ - Composted pressmud(12.5t ha⁻¹), T₄ - Farm yard manure (12.5t ha⁻¹), T₅ - NPK (100%), T₆ - Composted coirpith (12.5t ha⁻¹) + 50% NPK, T₇ - Composted pressmud(12.5t ha⁻¹) + 50% NPK, T₈ - Farm yard manure (12.5t ha⁻¹) + 50% NPK, T₉ - Composted coirpith (12.5t ha⁻¹) + 25% NPK, T₁₀ - Composted pressmud (12.5t ha⁻¹) + 25% NPK, T₁₁ - Farm yard manure (12.5t ha⁻¹) + 25 % NPK, T₁₂ - Composted coir pith (6.5t ha⁻¹) + Composted pressmud (6.5t ha⁻¹) + Farm yard manure (6.5t ha⁻¹)

Statistical analysis :

The data obtained on 25 DAS, 45 DAS, 55DAS and 75 DAS were subjected to the statistical analysis and based on the results inference were drawn.

RESULTS AND DISCUSSION

Table results obtained from the present investigation are presented in Table 1, 2 and 3.

Effect of composted pressmud, composted coirpith and FYM on vegetative characters of green gram :

The treatments T₂(composted coirpith) showed

increased root length on 25 DAS (13.47cm), T₁₀ (composted pressmud + NPK) on 45DAS (16.67cm) and T₅ (NPK 100%) on 55DAS (18.44cm) when compared to the control (2.44cm, 5.67cm and 9.11cm) on 25, 45 and 55 days after sowing, respectively. This is accordance with results of Kumarimani-muthuVeeral (2008) in maize. The significant increase in shoot length was obtained in T₅ (NPK 100%) on 25th (37.89cm) and T₆ (composted coirpith + 50%NPK) on 45th(47.32cm) days after sowing, T₃ (composted pressmud) on 55DAS (51.67cm) when compared to the control (24.56cm, 18.67cm and 22.33cm) on 25, 45 and 55 days after sowing, respectively. As in the present study, an increase in the shoot length was also noted by Hossain and Ishimine (2007) in turmeric.

The number of leaves were highest in T₇(12.88), T₁₂ (12.77), T₁₁ (12.22) treatment than the control T₁(6.77). On 45DAS the number of leaves were increased in T₃ (20.77) treatment when compared to the control T₁(8.55).

On 25DAS the number of nodules were significantly increased in T₄ (0.88, 7.00) when compared to the control T₁ (0.11 and 0.22) on 25 and 45 days after sowing. On 55DAS the number of nodules was increased in T₃ (8.66) when compared to the control T₁ (1.44). Number of flowers was increased in T₈ (7.67) and T₉ (6.33) on 45DAS when compared to the control T₁ (1.00).

Number of fruits was increased in T₆ (5.67), T₁₀ (5.00) on 55 DAS when compared to the control T₁ (1.33). This result is in agreement with findings of Pagaria and Totawat (2007). Plant fresh weight was increased in T₁₀ (4.93g), T₁₂ (4.82g), T₁₁ (4.87g) on 25DAS, T₈ (11.62g) on 45DAS and T₁₂ (63.54g) on 55DAS than control T₁ (1.18g, 1.65g and 5.83g) on 25, 45 and 55 days after sowing, respectively. Plant dry weight was

Table 1 : Effect of composted pressmud, composted coir pith and farm yard manure on vegetative parameters of green gram

Treatments	Root length(cm)			Shoot length(cm)			Number of leaves			Number of nodules		
	25DAS	45DAS	55DAS	25DAS	45DAS	55DAS	25DAS	45DAS	55DAS	25DAS	45DAS	55DAS
T ₁	2.44	5.67	9.11	24.56	18.67	22.33	6.77	8.55	14.66	0.11	0.22	1.44
T ₂	13.47	10.67	12.78	30.67	34.44	36.78	7.00	12.66	8.33	0.33	0.33	2.88
T ₃	6.56	5.78	12.56	36.44	43.78	51.67	7.11	20.77	8.77	0.66	0.33	8.66
T ₄	11.38	13.40	11.67	36.44	35.56	42.44	9.33	10.66	10.66	0.88	7.00	1.66
T ₅	13.37	6.74	18.44	37.89	45.56	48.44	8.00	10.66	10.88	0.55	0.22	1.77
T ₆	9.40	11.44	9.39	28.00	47.32	43.67	9.66	10.55	11.55	0.66	0.22	1.77
T ₇	8.44	8.38	11.41	27.89	35.00	46.89	12.88	11.55	14.11	0.55	3.66	1.55
T ₈	10.44	14.56	9.78	25.00	44.40	45.22	10.88	10.44	11.11	0.44	1.44	3.00
T ₉	6.46	14.78	14.89	31.44	31.56	34.22	7.55	13.44	10.88	0.44	2.88	2.77
T ₁₀	10.34	16.67	14.27	32.89	46.78	42.89	10.44	11.00	13.11	0.44	1.55	3.88
T ₁₁	8.67	9.44	10.78	37.78	43.33	45.67	12.22	13.00	13.44	0.33	2.00	3.55
T ₁₂	9.56	10.33	17.11	36.67	46.89	38.44	12.77	11.55	11.11	0.55	0.55	3.66
S.E.±		0.24944			0.90865			0.34058			0.28706	
C.D.(0.05)		0.49750			1.81226			0.67927			0.57253	
C.D.(0.01)		0.66051**			2.40607**			0.90184**			0.76013**	

** Indicate significance of value at P=0.01, DAS – Days after sowing

significantly high in T₁₁ (2.04g), T₁₀ (2.00g) on 25DAS T₈ (4.56g) on 45DAS and T₁₂ (5.59g) on 55 DAS when compared to the control T₁ (0.47g, 0.36g, 0.45g). Similar result was given by Akhilandeshwari (2003) in green gram and soybean.

Effect of composted pressmud, composted coirpith and FYM on yield parameters of green gram :

Number of pods per plant was increased in T₁₁ (4.77) when compared to the control T₁ (0.61). The length of pods

was significantly increased in T₁₀ (6.27cm), T₂ (6.24cm) and T₈ (5.78cm) when compared to the control T₁ (0.62cm).

The weight of the pods increased significantly in T₁₀ (4.92g) when compared to the control T₁ (0.57g). Similar was noted by Balakrishnan *et al.* (2009) and Umamaheshwari (2008) in soybean and cluster bean.

The number of seed per pod was significantly higher in T₅ (8.22), T₁₀ (8.00) and T₁₁ (7.77) When compared to the control T₁ (2.77). The weight of the seeds per pod was increased in the

Table 2 : Effect of composted pressmud, composted coirpith and farm yard manure on vegetative parameters of green gram

Treatments	Number of flowers		Number of fruits		Fresh weight (g)		Dry weight (g)	
	45DAS	55DAS	25DAS	45DAS	55DAS	25DAS	45DAS	55DAS
T ₁	1.00	1.33	1.18	1.65	5.83	0.47	0.36	0.45
T ₂	1.67	4.00	2.39	4.93	23.14	0.58	2.68	1.83
T ₃	2.00	4.00	1.78	6.73	21.68	0.55	2.45	2.61
T ₄	3.67	2.00	1.34	2.70	26.63	0.82	0.98	2.35
T ₅	4.67	4.33	2.11	5.19	32.19	0.69	1.80	3.48
T ₆	3.67	5.67	2.01	4.53	22.23	0.59	1.37	2.15
T ₇	4.67	2.33	2.68	6.14	32.18	1.00	1.58	1.68
T ₈	7.67	4.00	2.65	11.62	22.23	0.81	4.56	2.76
T ₉	6.33	4.67	2.01	4.18	23.13	0.30	1.65	1.53
T ₁₀	2.00	5.00	4.93	6.66	29.58	2.00	1.67	2.51
T ₁₁	3.67	3.33	4.87	3.36	35.18	2.04	1.23	3.01
T ₁₂	3.33	2.67	4.82	5.08	63.54	1.82	1.97	5.59
S.E.±	0.4714	0.4907		0.11534			0.06942	
C.D.(0.05)	0.9729	1.0127		0.23004			0.13845	
C.D.(0.01)	1.3258 **	1.3800 **		0.30541 **			0.18382 **	

** Indicate significance of value at P=0.01, DAS – Days after sowing

Table 3: Effect of composted pressmud ,coirpith, farm yard manure on yield parameters of green gram (75th day)

Treatments	Number of pods/plant	Length of pods(cm)	Weight of pods(g)	Number of seeds/pod	Weight of seed/pod (g)	Pod's fresh weight	Pod's dry weight
T ₁	0.61	0.62	0.57	2.77	0.03	0.05	0.11
T ₂	2.74	6.24	2.25	7.22	0.57	1.35	0.65
T ₃	3.65	5.37	2.38	3.77	1.33	3.26	0.14
T ₄	3.73	4.78	2.40	4.33	0.41	2.26	0.06
T ₅	3.61	5.68	3.12	8.22	2.13	3.12	0.46
T ₆	2.78	5.26	0.67	3.11	0.22	0.54	0.55
T ₇	2.67	5.32	2.67	7.11	1.42	2.67	0.12
T ₈	2.72	5.78	3.56	6.66	4.37	3.57	0.22
T ₉	1.68	5.21	1.61	5.33	0.20	1.62	0.37
T ₁₀	2.76	6.27	4.92	8.00	3.46	4.93	0.03
T ₁₁	4.77	5.75	3.37	7.77	1.77	3.34	0.12
T ₁₂	3.27	5.77	2.72	6.09	1.59	2.71	0.25
S.E.±	0.0779	0.0706	0.0602	0.5379	0.8626	0.2419	0.0470
C.D.(0.05)	0.1617	0.1465	0.1248	1.1156	1.7889	0.5016	0.0974
C.D.(0.01)	0.2197 **	0.1991 **	0.1696 **	1.5164 **	2.4315 **	0.6818 **	0.1324**

** Indicate significance of value at P=0.01, DAS – Days after sowing

T₈(4.37g) when compared to the control T₁ (0.03g).

The pods fresh weight was increased in T₁₀(4.93g) when compared to the control T₁(0.05g).The results were in agreement with the findings of Chandrasekaran *et al.* (2007) in groundnut. The pods dry weight was highest in T₂ (0.65g) when compared to the control T₁(0.11g). This is accordance with result of Muthuraju and Siddaranappa (2005).

Conclusion :

Disposal of waste material into the environment causes pollution, health hazards to the people and affects the crop productivity. Organic manures improve the physical, chemical and biological properties of the soil.

From the present investigation it has become evident that the composted pressmud, composted coirpith and FYM along with NPK influence the biometrical and yield characters of green gram. Hence it was concluded that the coirpith and pressmud on composting can be used as an organic manure substitute.

REFERENCES

- Akilandeswari, A. (2003). Impact of composted pressmud on seed germination, seedling growth, biometric and yield parameters of green gram and soyabean, M.Sc. Thesis, Avinashilingam Deemed University, Coimbatore, T.N. (INDIA).
- Balakrishnan, V., Ravindran, K.C., Snajiviraja, K. and Venkatesan, K. (2009). Halopahytic compost for sustainable agriculture. *Science*, **5**(3) : 56 – 59.
- Chandrasekaran, R., Somasundaram, E., Mohammad Amanullah, Nalini, K., Thirukkumaran, K. and Sathyamoorthy, K. (2007). Response of confectionery groundnut (*Arachis hypogea* L.) varieties to farm yard manure. *J. Appl. Sci. Res.*, **3**(10) : 1097–1099.
- Hossain, Md.Amzad and Ishmine, Yukio (2007). Effect of FYM on growth and yield of turmeric (*Curcuma longa* L.) cultivated in Dark – Red soil, Red soil and Grey soil in Okinawa, Japan. *Plants prod. Sci.*, **10** (1):146 – 150.
- Kumarimanimuthu Veeral, D. (2008). Utilization of agro industrial waste in maize based cropping system, *Agric.Sci.Digest.*, **28** (3):195 – 197.
- Muthuraju, M., Ravi, M.V. and Siddaramappa, R. (2005). Effect of application of enriched pressmud on changes in chemical properties of an alfisol. *Mysore Agric. Sci.*, **39**(2) : 207 – 213.
- Pagaria, P. and Totawat, K.L. (2007).Effect of pressmud and spent wash in integration with phosphogypsum on metallic cation build – up in the calcareous sodic soil. *J. Indian Soc. Soil Sci.*, **55** (1) : 52 – 57.
- Sneha, Raut, S., Chare, C.N., Deotale, R.D., Waghmare, H.V., Hatmode, C.H. and Yen Prediwar, N.D.(2003). Response of seed dressing with biofertilizers and nutrient on morpho – physiological parameters and yield of soyabean. *J. Soils & Crop*, **13**(2) : 309 – 313.
- Umamaheshwari, P.(2008). Influence of composted pressmud on the soil health and productivity of legumes *Cyamopsis tetragonoloba* L. Var. Pusa Navababar and [*Glycine max* (L.)(Merrill Var.Co.Soy3)], M.Sc., Thesis, Avinashilingam Deemed University, Coimbatore, T.N. (INDIA).

8th
Year
★★★★★ of Excellence ★★★★★