



## Effect of potassium and zinc on yield, quality and economics of Gujarat cowpea-4

A.S. CHAVAN\* AND H.R.KHAFI<sup>1</sup>

Department of Agronomy, Marathwada Krishi Vidyapeeth,  
PARBHANI (M.S.) INDIA (Email : anil.bunty212@gmail.com)

**Abstract :** A field experiment was conducted on medium black calcareous soil of the Instructional Farm, Junagadh Agricultural University, Junagadh (Gujarat) during the season of *Kharif*-2008. Significantly higher grain yields ( $1587 \text{ kg ha}^{-1}$ ) were recorded with application of  $60 \text{ kg K}_2\text{O ha}^{-1}$ . Significantly highest grain yields ( $1553 \text{ kg ha}^{-1}$ ) were recorded with  $40 \text{ kg zinc ha}^{-1}$ . The maximum grain protein content of 24.85 per cent as well as The higher net realization Rs.21725  $\text{ha}^{-1}$  with BCR (3.43) was accrued under the application of  $60 \text{ kg K}_2\text{O ha}^{-1}$ . The higher net realization of Rs.20545  $\text{ha}^{-1}$  with 3.18 BCR was realized under the application of  $40 \text{ kg zinc ha}^{-1}$ . Considering the treatment combinations,  $\text{K}_3\text{Zn}_2$  ( $60 \text{ kg K}_2\text{O ha}^{-1} + 40 \text{ kg zinc ha}^{-1}$ .) recorded the highest net realization of Rs. 24344  $\text{ha}^{-1}$  with BCR 3.54. It can be indicated that the potential production and profit from *Kharif* season cowpea (cultivar GC-4) can be secured by fertilizing the crop with  $60 \text{ kg K}_2\text{O ha}^{-1}$  along with  $40 \text{ kg zinc ha}^{-1}$ .

**Key Words :** Cowpea, Potash, Zinc

**View Point Article :** Chavan, A.S. and Khafi, H.R. (2013). Effect of potassium and zinc on yield, quality and economics of Gujarat cowpea-4 . *Internat. J. agric. Sci.*, 9(1): 329-331.

**Article History :** Received : 17.11.2011; Revised : 03.12.2012; Accepted : 27.12.2012

### INTRODUCTION

Cowpea (*Vigna sinensis* L.) is the most versatile *Kharif* as well as summer pulse, because of its smothering nature, drought tolerant character, soil restoring properties and multipurpose uses. It covers the ground and checks soil erosion and works as mulch to reduce the evaporation losses apart from being a leguminous crop. Cowpea can fix about 80 to 90  $\text{kg N ha}^{-1}$  under ideal condition. At present India is passing through a shortage of protein where the people are predominantly vegetarian, pulses are the main source of protein and thus are of vital importance in daily diet. However, requirement of pulses are going up due to population explosion while its production is not increasing to that extent consequently, the price of pulses has increased exorbitantly and common man can not afford to purchase the same. Hence, the production of pulses have to be increased either by increasing the land or by increasing productivity.

Amongst the nutrients N and P are given the priority and very little attention is paid towards the K and micronutrients which are of prime importance for the nutrition of cowpea from the nutrition point of view.

The potassium is one of the major plant nutrient for the growth and development of plants. The major functions are associated with enzyme involved in photosynthesis, metabolism of carbohydrate and physiological processes, such as root growth, water uptake and utilization efficiency, synthesis of protein and amino acids, enzyme activation and yield determining process *viz.*, drought, pest and disease tolerance.

Zinc plays vital role in plant growth and development. Zinc also catalyses the biosynthesis of indole acetic acid (IAA), acting as metal activator of the enzyme, there by ultimately increasing crop yield. Moreover, it controls the equilibrium between  $\text{CO}_2$ , water and carbonic acid in plant metabolism and helps in synthesis of nucleic acids, proteins and stimulates seed formation. Its deficiency retards

\* Author for correspondence

<sup>1</sup>Pearl Millet Research Station, JAMNAGAR (GUJARAT) INDIA

photosynthesis and nitrogen metabolism.

Thus, arriving at an optimum combination of potassium and zinc chemical fertilizers would result in boosting the production of cowpea, particularly that of newly developed variety, Gujarat Cowpea-4.

With the background and inspite of adequate evidence on its productive capabilities the information on agro-techniques for cowpea production, especially its nutrient requirements for medium black soil of Saurashtra region in *Kharif* season is lacking. A proper understanding of the nutrient management of cowpea crop especially in *Kharif* season is an urgent need to enhance production. So far there is no practically systematic research work done to evaluate the effect of potassium and zinc on growth, yield attributes, quality and nutrient uptake of cowpea crop in this region.

## MATERIALS AND METHODS

A field experiment was conducted during the *Kharif* season 2008 at Agronomy farm, college of Agriculture, Junagadh Agricultural University, Junagadh (Gujarat). The experiment comprising of twelve treatment combinations consisting of four levels of potassium *viz.*, no potassium application ( $K_0$ ), 20 kg  $K_2O$  ha<sup>-1</sup> ( $K_1$ ), 40 kg  $K_2O$  ha<sup>-1</sup> ( $K_2$ ), 60 kg  $K_2O$  ha<sup>-1</sup> ( $K_3$ ) and three

levels of zinc *viz.*, no zinc application ( $Zn_0$ ), 20 kg zinc ha<sup>-1</sup> ( $Zn_1$ ) and 40 kg zinc ha<sup>-1</sup> ( $Zn_2$ ) were framed in Factorial Randomized Block Design (FRBD) with four replications. The soil experimental site was on medium black calcareous soil with pH of 7.9 which was free from any kind of salinity or sodicity hazards. The gross and net plot sizes were 5.00 m x 3.6 m and 4.00 m x 2.4 m, respectively. Potash in the form of murate of potash (60%  $K_2O$ ) and zinc in the form of zinc sulphate (21% Zn) was applied at the time of sowing in furrows as per treatments.

## RESULTS AND DISCUSSION

The results of the present study as well as relevant discussions have been presented under following sub heads:

### Effect of potash :

A perusal of data given in Table 1 revealed that the grain yield was significantly influenced by potassium levels. Significantly the highest grain yield of 1587 kg ha<sup>-1</sup> was recorded with 60 kg  $K_2O$  ha<sup>-1</sup> as compared to  $K_2$  (40 kg  $K_2O$  ha<sup>-1</sup>),  $K_1$  (20 kg  $K_2O$  ha<sup>-1</sup>) and  $K_0$  (no potassium application) which recorded grain yield 1454, 1313 and 1003 kg ha<sup>-1</sup>, respectively. The results are in agreement with findings of Cheng *et al.* (1999), Gill and Kamprath (1990), Patra *et al.* (1995), Ghatak *et*

**Table 1: Effect of different treatments on grain and protein content of cowpea**

Treatments	Grain yield (kg ha <sup>-1</sup> )	Protein content (%)
Potassium (K)		
$K_0$ (control)	1003	19.31
$K_1$ (20 kg ha <sup>-1</sup> )	1313	22.54
$K_2$ (40 kg ha <sup>-1</sup> )	1454	23.82
$K_3$ (60 kg ha <sup>-1</sup> )	1587	24.85
S.E. ±	45.50	0.65
C.D. at 5%	131.01	1.86
Zinc (ZnSO <sub>4</sub> )		
$Zn_0$ (control)	1024	21.05
$Zn_1$ (20 kg ha <sup>-1</sup> )	1441	23.26
$Zn_2$ (40 kg ha <sup>-1</sup> )	1553	23.59
S.E. ±	39.40	0.56
C.D. at 5%	113.46	1.61
C.V. %	11.77	9.88
Interaction-K X Zn	Sig.	NS

**Table 2: Effect of different treatments on GMR, NMR and B:C ratio of cowpea**

Treatments	Gross realization (Rs. ha <sup>-1</sup> )	Net realization (Rs. ha <sup>-1</sup> )	Benefit: cost ratio
Potassium (K)			
$K_0$ (control)	19395	10816	2.218
$K_1$ (20 kg ha <sup>-1</sup> )	25360	16677	2.908
$K_2$ (40 kg ha <sup>-1</sup> )	28051	19262	3.184
$K_3$ (60 kg ha <sup>-1</sup> )	30616	21725	3.430
Zinc (ZnSO <sub>4</sub> )			
$Zn_0$ (control)	19755	11698	2.443
$Zn_1$ (20 kg ha <sup>-1</sup> )	27853	19117	3.183
$Zn_2$ (40 kg ha <sup>-1</sup> )	29959	20545	3.180

al. (1997) and Swaroop and Rathore (2002).

An appraisal of data presented in Table 1 showed that significantly higher protein content of 24.85 per cent was recorded when crop was fertilized with 60 kg K<sub>2</sub>O ha<sup>-1</sup> (K<sub>3</sub>), which remained at par with 20 kg K<sub>2</sub>O ha<sup>-1</sup> (K<sub>1</sub>), and no potassium application (K<sub>0</sub>) which recorded 22.54 and 19.31 per cent, respectively. The results are in accordance with those reported by Nanjundappa and Manure (2002), Bansal *et al.* (2001).

There was considerable increase in net realization due to potassium application as given in Table 2. The highest net return of Rs.21725 was obtained with 60 kg K<sub>2</sub>O ha<sup>-1</sup> (K<sub>3</sub>) followed by Rs.19262 with (K<sub>2</sub>) 40 kg K<sub>2</sub>O ha<sup>-1</sup> and Rs.16677 with 20 kg K<sub>2</sub>O ha<sup>-1</sup> (K<sub>1</sub>). The lowest net realization of Rs.1086 ha<sup>-1</sup> was noted with K<sub>0</sub> (no potassium application). The results are in accordance with those reported by Nanjundappa and Manure (2002), Bansal *et al.* (2001).

#### Effect of zinc :

An appraisal of results in Table 1 showed that the effect of zinc on grain yield was found significant. Significantly the highest grain yield by 1553 kg ha<sup>-1</sup> was observed with application of zinc 40 kg ha<sup>-1</sup> (Zn<sub>2</sub>) as compared to 20 kg zinc ha<sup>-1</sup> (Zn<sub>1</sub>) and Zn<sub>0</sub> (no zinc application). The grain yield obtained under Zn<sub>1</sub> and Zn<sub>0</sub> were 1441 and 1024, respectively. The results are in agreement with those reported by Vedram *et al.* (2002), Patil *et al.* (2006), Jain and Dahama (2006), Sakal (2001), Dadhich and Gupta, (2005), Subramaniyan *et al.* (2002), Choudhary *et al.* (2005), Deosarkar *et al.* (2001).

The results in Table 1 indicated that protein content in grain was significantly affected due to varying zinc levels. Significantly higher protein content 23.59 per cent was noted under zinc 40 kg ha<sup>-1</sup> (Zn<sub>2</sub>), which remained statistically at par with 20 kg zinc ha<sup>-1</sup> (Zn<sub>1</sub>), which recorded 23.26 per cent protein content. Significantly the lowest protein content 21.05 per cent was recorded under Zn<sub>0</sub> (no zinc application). The present findings are in agreement with those reported by Vedram *et al.* (2002), Shelge *et al.* (2000) and Jain and Dahama (2006).

There was considerable increase in net realization due to potassium application as given in Table 2. The highest net return of Rs.21725 was accrued with 60 kg K<sub>2</sub>O ha<sup>-1</sup> (K<sub>3</sub>) followed by Rs. 19262 with (K<sub>2</sub>) 40 kg K<sub>2</sub>O ha<sup>-1</sup> and Rs.16677 with 20 kg K<sub>2</sub>O ha<sup>-1</sup> (K<sub>1</sub>). The lowest net realization of Rs.10816 ha<sup>-1</sup> was realized with K<sub>0</sub> (no potassium application).

The economics presented in Table 2 revealed that there was a considerable increase in net realization up to Zn<sub>2</sub> level (40 kg ZnSO<sub>4</sub> ha<sup>-1</sup>). The maximum net realization of Rs. 20545 ha<sup>-1</sup> was obtained with Zn<sub>2</sub> level, which was higher than Rs.19117 ha<sup>-1</sup> and Rs.11698 ha<sup>-1</sup>, obtained in 20 kg ZnSO<sub>4</sub> ha<sup>-1</sup> (Zn<sub>1</sub>) and no zinc application (Zn<sub>0</sub>), respectively. The results are in accordance with those reported by Nanjundappa and Manure (2002), Bansal *et al.* (2001).

## REFERENCES

- Bansal, K.N., Singh, S.P. and Nepalia, V. (2001).** The effect of potassium application on yield and quality of soybean and wheat in Madhya Pradesh. *Fertiliser News*, **46**(11): 45-48 & 51-52.
- Cheng, G. H., Li, C. J., Jiang, G. and Zhao, G. Z. (1999).** Influence of main nutrient elements on soybean yield. *J. Jilin Agric. Univ.*, **21**(3): 68-70.
- Choudhry, B. R., Keshwa, G. L. and Parihar, C. M. (2005).** Effect of thiourea and zinc on productivity of pearl millet. *Ann. Agric. Res.*, **26** (3): 424-427.
- Dadhich, L.K. and Gupta, A.K. (2005).** Growth and yield of fodder pearl millet as influenced by sulphur, zinc and intercropping with cowpea. *Fertilizer News*, **50**(3): 55-57.
- Deosarkar, D.B., Patinge, S.P., Bhosale, A. M. and Deshmukh, S.B. (2001).** Effect of micronutrients on seed yield of soybean. *Ann. Plant Physiol.*, **15**(2): 163-166.
- Ghatak, S., Sounoa, G., Maitra, S., Saron, B.K., Roy, D.K. and Panda, P.K. (1997).** Effect of irrigation and potassium on growth and productivity of summer groundnut. *Environ. & Ecol.*, **15**(2): 469-473.
- Gill, D.W. and Kamprath, E.J. (1990).** Potassium uptake and recovery by an upland rice-soybean rotation on an Oxisol. *Indian J. Agron.*, **82**(2): 329-333.
- Jain, N.K. and Dahama, A.K. (2006).** Direct and residual effect of P and Zn fertilization on productivity of wheat-pearlmillet cropping system. *Indian J. Agron.*, **51**(3): 165-169.
- Nanjundappa, G. and Manure, G.R. (2002).** Influence of phosphorus and potassium levels on yield and quality of fodder cowpea [*Vigna unguiculata* (L.) Walp.]. *Mysore J. Agric. Sci.*, **36**(2): 104-106.
- Patra, A.K., Tripathi, S. K., Samui, R. C. (1995).** Seasonal variation in growth and yield of groundnut at different levels of N and K. *Indian J. Plant Physiol.*, **38** (3): 218-223.
- Patil, S.R., Gitte, A. N., Shelke, B.M. and Tike, M.A. (2006).** Influence of Zn and B on yield and yield contributing characters in sunflowers. *J. Maharashtra Agric. Univ.*, **31**(3): 247-208.
- Sakal, R. (2001).** Efficient management of micronutrients for sustainable crop production. *J. Indian Soc. Soil Sci.*, **49**(4): 593-608.
- Shelge, B. S., Sontakke, J. S. and Sondge, V. D. (2000).** Influence of micronutrients on yield of soybean. *Madras Agric. J.*, **87**(7-9): 538-540.
- Subramaniyan, K., Kaliselvan, P. and Arulmozhi, N. (2002).** Response of confectionary groundnut to micronutrients. *Legume Res.*, **24**(2): 139-140.
- Swaroop, Kishan and Rathore, S.V.S (2002).** Economics, nutrient content and pod yield of vegetable cowpea in relation to application of P K and *Rhizobium* biofertilizer in Andaman. *Indian-Agriculturist*, **46** (3/4): 153-160.
- Vedram Misra, S.K. and Upadhyay, R.M. (2002).** Effect of S, Zn and biofertilizers on quality characters in mungbean. *Indian J. Pulse Res.*, **15** (2) : 139-141.

\*\_\*\_\*\_\*\_\*\_\*\_\*\_\*