

RESEARCH PAPER

DOI: 10.15740/HAS/IJPPHT/9.2/44-48

Temperature dependant hydration kinetics of blanched and unblanched Amarnath (Rajgira) leaves

■ Virendra Foke* and Dhananjay Kulkarni

Department of Agricultural Processing and Food Engineering, Shriram College of Agricultural Engineering, Paniv, Solapur (M.S.)
India

Email : virendrafoke@gmail.com; dhanukulkarni@gmail.com

*Author for Correspondence

■ Research chronicle : Received : 20.10.2018; Revised : 20.11.2018; Accepted : 23.11.2018

SUMMARY :

A study was conducted to determine the effect of blanching on moisture adsorption and desorption characteristics of amaranth leaves by static gravimetric method. Different saturated salt solutions were prepared to provide constant relative humidity environments. Four amaranth leaves samples (Unblanched, Unblanched dried, Blanched and Blanched dried) were kept at 3 temperatures (30°C, 40°C and 60°C) with corresponding five relative humidity levels (09.76 to 90.28 %). It was observed that EMC of leaves increased with increase in relative humidity at constant temperature also at constant ERH, EMC decreased with increase in temperature. The adsorption EMC was lower than the desorption EMC for all the samples at all temperature and relative humidity ranges. The blanched dried amaranth leaves exhibited lowest EMC than other samples. Sorption data was modeled by using Henderson equation to develop guideline for moisture sorption process. Both constant of equation found to be dependent on temperature. The relation between constant was expressed by a polynomial equation.

KEY WORDS : Temperature dependant, Hydration kinetics, Blanched, Unblanched Amarnath leaves

How to cite this paper : Foke, Virendra and Kulkarni, Dhananjay (2018). Temperature dependant hydration kinetics of blanched and unblanched Amarnath (Rajgira) leaves. *Internat. J. Proc. & Post Harvest Technol.*, **9** (2) : 44-48. DOI: 10.15740/HAS/IJPPHT/9.2/44-48. Copyright@ 2018: Hind Agri-Horticultural Society.

A marnath (Rajgira) leaves (*Amranthus*) is an important leafy vegetable commonly grown in India. It contains 92.10 per cent moisture, 2.2 per cent carbohydrate, 3.4 per cent protein, 2.2 per cent fibre, 0.4 per cent fat and rich source of vitamin A, iron and calcium (Anonymous, 2009). Shelf-life of rajgira leaves is shorter due to its high moisture content.

Therefore, to increase shelf-life, its moisture should

be removed which can be done by dehydration and stored at particular temperature and relative humidity. However, changes in flavour and textural alterations may occur during storage of dried food products. These changes are particularly influenced by moisture content of material, relative humidity and storage conditions. Hence, knowledge of such transformation factors is important for predicting the various storage conditions of particular

material (Velez-Ruiz *et al.*, 2004).

EMC is defined as moisture content of hygroscopic material in equilibrium with a particular environment in terms of temperature and relative humidity. The corresponding relative humidity is known as equilibrium relative humidity (ERH). Material can attain EMC either by losing the moisture (desorption isotherm) or by gaining the moisture from surrounding (adsorption isotherm). Thus with the knowledge of the moisture sorption isotherms of vegetable plant, it is possible to predict the maximum moisture that the plant can be allowed to gain during storage. Sorption isotherms are an extremely valuable tool for scientists because they can be used to predict potential changes in biological material's stability. The adsorption isotherm data can be used for a storage method determination while the desorption isotherm data can be used for drying analysis (Daniel and Wanderley, 2004).

Mathematical modeling of sorption process is known to be important for the design of processing equipments and optimization of storage and drying conditions. Work has been conducted on the study of sorption isotherms of different medicinal plants like tow mint, aroma leaves, calendula leaves and so on. (Kouhila *et al.*, 2007; Silva *et al.*, 2004 and Velez-Ruiz *et al.*, 2004). However, sufficient information about vegetable like amaranth leaves appears to be lacking. A study was, therefore, undertaken for blanched and unblanched rajgira leaves. This information will be useful in identifying the most shelf-stable product of it and appropriate packaging materials.

EXPERIMENTAL METHODS

About 10 kg of fresh rajgira leaves was purchased from a local market. Air tight glass containers of 350 ml capacity each, five analytical salts namely Lithium chloride (LiCl), magnesium chloride (MgCl₂), sodium bromide (NaBr), sodium chloride (NaCl) and potassium nitrate (KNO₃) with purity 99.9 per cent were used.

Sample preparation:

Fresh rajgira leaves were thoroughly washed and rinsed with potable water. The washed leaves were randomly selected, divided into two portions and cut into irregular shapes of about 1 cm³ size. One portion was used as it is *i.e.* as unblanched while second portion was blanched by dipping the sample in boiling water (98±2°C) for two min. and then immediately cooling in water

(Chakraverty and De, 1981). The half part of each of unblanched and blanched portion was dried in cabinet dryer at 60°C for 4 hr to get the four samples namely unblanched rajgira, unblanched dried rajgira, blanched rajgira and blanched dried rajgira leaves. After determining moisture content (% d.b.) of all these samples by oven method (105°C for 24 h), they were used for sorption study. Unblanched and blanched rajgira leaf portions were used for desorption study while unblanched dried and blanched dried rajgira portions were used for adsorption study.

Determination of sorption isotherms:

The setup for sorption study consists of air tight glass container with metal lid equipped with saturated salt solution and air oven. For sorption study of samples, saturated salt solutions were prepared (Silva *et al.*, 2004 and Zomorodian and Tavakoli, 2007), which gave relative humidity in the range of 09.76 to 90.28 per cent at constant temperatures (Table 1). Each of 100ml of above prepared saturated salt solution was filled in different glass containers and kept at constant temperatures (30, 40 and 50°C) in oven for 24 h for obtaining constant relative humidity (Kouhila *et al.*, 2007). After completion of equilibration process, each samples of four type of above prepared leaves (1 g each for desorption and adsorption study) were kept in glass container holder. After hermitically sealing the containers, they were again kept in oven for EMC determination till constant weight of samples was reached (Alakali *et al.*, 2009). Each sample had taken around eight hours to reach constant weight. Each experiment was replicated thrice. After obtaining constant weight of each sample, its EMC (% d.b) was determined by using standard air oven method at 105°C for 24 h (AOAC, 1990).

Modelling of sorption isotherms:

Henderson equation was used to model the sorption isotherm for all the samples of spinach as shown in following eq.

$$1 - ERH = e^{-cTM^n} \quad \dots(1)$$

Here, ERH is equilibrium relative humidity (decimal), Me is equilibrium moisture content (% d.b.), T is temperature (°C) and c and n are constant depend on crop type.

The eq. (1) was rearranged in the form of eq. (2)

and (3),

$$\log(1 - ERH) = -cTm_e^n \dots(2)$$

$$\log\left(\frac{-\log(1 - ERH)}{T}\right) = \log c + n \log M_e \dots(3)$$

It was considered that linear regression analysis (Microsoft excel 2007) of $\left(\frac{-\log(1 - ERH)}{T}\right)$ against $\log M_e$ gives value of co-efficients (c and n) of Henderson equation.

EXPERIMENTAL FINDINGS AND ANALYSIS

The initial moisture contents of unblanched, blanched, unblanched dried and blanched dried rajgira leaves were observed, 89.30, 90.79, 3.37 and 2.18 per cent (w.b.), respectively. Equilibrium moisture contents (EMC) with equilibrium relative humidity (ERH) for all the types of leaves shown in Table 1. It can be seen that EMC of

sample increased with increase in ERH at a given temperature. Also it was observed that at a particular ERH, EMC of samples decreased with increase in temperature. This might be due to rajgira leave samples became less hygroscopic with increase in temperature at constant relative humidity. Similar trends were observed for maytenus leaves (Daniel and Wanderley, 2004); tow mint (Kouhila *et al.*, 2007) and calendula leaves (Silva *et al.*, 2004).

EMC of blanched rajgira sample always exhibited less EMC than unblanched sample at that particular relative humidity (Table 1). This might be due to availability of higher moisture in the container at higher relative humidity levels. These observations are in line with the findings of Velez-Ruiz *et al.* (2004); Kouhila *et al.* (2007) and Alakali *et al.* (2009). Blanched and blanched dried samples gave lower EMC as compared to unblanched and unblanched dried leaves, respectively. The EMC's of blanched leaves were lower than unblanched, because

Table 1: Desorption and adsorption equilibrium moisture contents of rajgira leaves at different temperature and relative humidity levels

Temperature (°C)	Equilibrium relative humidity (ERH) (%)	Desorption EMC (% d.b)		Adsorption EMC (% d.b)	
		Unblanched	Blanched	Unblanched	Blanched
30	10.22	7.58	6.14	4.53	4.12
	31.12	10.91	9.87	10.02	9.68
	53.60	19.92	16.76	17.03	14.16
	77.14	27.85	24.33	22.89	20.95
	90.28	44.74	42.88	40.87	40.21
40	10.19	6.62	4.91	4.19	4.05
	30.78	9.91	9.78	9.31	7.78
	53.20	19.11	15.88	15.01	14.32
	73.68	27.46	22.25	20.68	18.35
	88.23	42.80	41.56	40.19	36.46
60	09.76	4.43	3.77	3.06	3.07
	27.42	8.94	7.39	6.23	5.12
	45.23	16.03	13.57	12.49	10.35
	71.40	23.69	20.72	18.55	15.47
	84.10	38.40	35.89	34.04	32.41

Table 2 : Henderson equation constants for Desorption and adsorption isotherms for unblanched rajgira leaves

Temperature (°C)	Henderson equation constants unblanched spinach leaves					
	Desorption isotherm			Adsorption isotherm		
	c	n	R ²	c	n	R ²
30	0.0262	1.328	0.998	0.0287	1.342	0.990
40	0.0260	1.224	0.993	0.0310	1.252	0.960
50	0.0232	1.230	0.982	0.0325	1.047	0.970

Table 3: Henderson equation constants for Desorption and adsorption isotherms for blanched rajgira leaves

Temperature (°C)	Henderson equation constants for blanched spinach leaves					
	Desorption isotherm			Adsorption isotherm		
	c	n	R ²	c	n	R ²
30	0.0306	1.160	0.984	0.0326	1.415	0.977
40	0.0270	1.101	0.989	0.0347	1.183	0.969
50	0.0268	1.090	0.978	0.0389	0.982	0.986

Table 4: Relation between Henderson equation constants for Desorption and adsorption isotherms for rajgira leaves

Desorption isotherm (unblanched leave)	$c = -8x 10^{-6}T^2 + 0.0006T + 0.0141$ (R ² = 1) $n = 0.0003T^2 - 0.0344T + 2.078$ (R ² = 1)
Adsorption isotherm (unblanched dried)	$c = -1.4x 10^{-6}T^2 + 0.0002T + 0.0186$ (R ² = 1) $n = 0.0002T^2 - 0.0268T + 1.75$ (R ² = 1)
Desorption isotherm (blanched leave)	$c = 8.5x 10^{-6}T^2 - 0.0007T + 0.03$ (R ² = 1) $n = -5.8x 10^{-5}T^2 + 0.001T + 1.32$ (R ² = 1)
Adsorption isotherm (blanched dried)	$c = 2.7x 10^{-6}T^2 - 0.0001T + 0.024$ (R ² = 1) $n = 0.0003T^2 - 0.033T + 1.69$ (R ² = 1)

of its less hygroscopic nature, which may be due to the hot water blanching which lead to gelatinization of chlorophyll in the leaves possibly creating a less hygroscopic polymer network around the blanched dried surfaces. The same trend was observed for ginger by Alakali *et al.* (2009).

Henderson equation constants for desorption and adsorption isotherms for unblanched and blanched rajgira leaves shown in Table 2 and 3. It can be seen that co-efficient of determination was higher (R² > 0.95) indicated that Henderson equation satisfactorily described sorption characteristics for all the samples. The constant 'c' was almost equal indicated that there was no much effect of temperature on sorption process but there might be possibility of higher variation with higher temperature variation in sorption process. The constant 'n' was decreased with increase in temperature for both adsorption and desorption process for all the samples. This effect might be depending on moisture transfer or moisture diffusion properties of leaf (Silva *et al.*, 2004). Therefore, relation between these co-efficients with temperature was expressed by using polynomial equation (R²=1) as shown in Table 4.

Conclusion:

EMC of amaranth (Rajgira) leaves increased with increase in relative humidity at all temperatures, while it decreased with increase in temperature at all relative

humidity levels. Also, blanching greatly affected the adsorption and desorption characteristics of rajgira leaves. Blanched dried samples exhibits lowest EMC at all temperature and relative humidity ranges. Henderson equation satisfactorily (R² > 0.95) described moisture sorption characteristics for all the samples of rajgira leaves.

LITERATURE CITED

- Alakali, J. S., Irtwange, S. V. and Abu, J. O. (2009).** Effect of processing methods and storage environment on moisture adsorption characteristics of ginger. *African J. Food, Agric., Nutr. & Develop.*, **9**(5): 1245-1257.
- Anonymous (2009). USDA National Nutrient database for standard Reference, release 22, U.S, Department of Agriculture, Mary land.
- AOAC (1990). Moisture content in plants. *Association of Official Analytical Chemists*, **1**: 949.
- Chakraverty, A. and De (1981).** *Post harvest technology of cereals, pulses and oilseeds*. IBH Publication. 54pp.
- Daniel, S. and Wanderley, P. O. (2004).** Drying of medicinal plants. Equilibrium moisture content and mathematical modeling of maytenus ilicifolia leaves. *Proceedings of the 14th International Drying Symposium*, **C**: 1712-1719.
- Kouhila, M., Ethmane, C. S., Lamharrar, A., Idlimam, A. and Mimet, A. (2007).** Moisture sorption isotherms and thermodynamic properties of tow mints: Mentha pulegium and mentha rotundifolia. *Revue des Energies*

Renouvelables, **11**(2): 181-195.

Silva, F. D., Park, K., Magalhaes, P. and Pozitano, M. (2004).
Desorption isotherms of calendula officinalis leaves.
Proceedings of the 14th International Drying Symposium,
C: 1569-1576.

Velez-Ruiz, J. F., Lima, M. E. and Macedo, R. C. (2004). Air

drying of three aroma herbs Basil, Dill and Tea herbs.
Proceedings of the 14th International Drying Symposium,
C: 1914-1921.

Zomorodian, A. A. and Tavakoli, R. A. (2007). The adsorption-
desorption hysteresis effect on pistachio nuts. *J. Agric.*
Sci. Technol., **9**: 259-265.

9th
Year
★★★★★ of Excellence ★★★★★